



PRODUCTIVITY AND OIL QUALITY OF *THYMUS VULGARIS* L. UNDER GROWTH PROMOTERS AND SOILLESS CULTURE CONDITIONS

Fatma S. Aboud, M.H. Mohammed, Nagwa M. Ahmed and S.H. Ahmed

Closed Agricultural Methods Research Dept., Central Laboratory for Agriculture Climate, Agriculture Research Center.

Abstract

Thyme (*Thymus vulgaris* L.) belongs to the lamiaceae family. It is characterized by its medicinal and economic values. Thymus is grown worldwide for cosmetic and medical use. Plant bio-stimulants are classified as substances which have positive effects on growth and productivity of plants. Bio-stimulants are obtained naturally from various economically and environmentally viable sources. The current experiment aimed to investigate the response of thymus physical and chemical properties to the different types and concentrations of bio-stimulators under soilless culture conditions for two successive seasons 2019 and 2020. Thymus seedlings were planted in pots filled with mixed media of perlite: sand (1:1 v/v). Plants were sprayed with aqueous solution of ascorbic acid (1 and 2g/l), aspartic acids (500 and 1000 ppm) and dry yeast (2 and 4g/l) and combination of them compared with control. The herb was harvested in mid- July (First cutting) and first week of October (second cut). Growth and yield characters were measured. The essential oil percentage was determined and analyzed by GC in both cuts from the fresh herb. Results showed that, spraying of mixtures of ascorbic acid (2g/l), aspartic acids (1000 ppm) and yeast (4g/l) significantly increased yield, total carbohydrates, total phenols, total pigments, essential oil percentage and yield as well as its NPK content characters and components of essential oil high contents of thymol, α and β - Pinene during both cuts and for both seasons. Economic evaluation was carried out by calculating with the highest return in the mixture treatment. It is clear that all treatments are economically feasible however, one can find that the maximum return or profit comes from mixture treatments where the benefit to cost ratio were 3.99 and 3.9 for Mix2 and Mix1 respectively.

Key words: *Thymus vulgaris* L., substrate Culture, vitamins, amino acids, active dry yeast, thymol, economic evaluation.

Introduction

Medicinal plants have an important value due to their natural antioxidant and antimicrobial activities, of them is thymus. Many authors confirmed the medicinal and economic importance of thymus (Satyal *et al.*, 2016; Aouam *et al.*, 2019; Golkar *et al.*, 2020) where it can be consumed locally or exported. Medicinal plants have an important socio-cultural, spiritual and medicinal value in rural and tribal lives (Shinwari, 2005) of them, Thymus plants which belongs to the Lamiaceae family and is characterized by its high nutritional and medicinal value. Thymus is rich in nutrients, thiamine, riboflavin and niacin. Thymus has been reported for its antioxidative, antimicrobial, antitussive expectorant, antispasmodic and antibacterial effects (Sarikurku *et al.*, 2015; Hashem 2018). Thymus essential oil contains thymol, carvacrol, linalool, α -terpineol, camphor, caryophyllene and γ -terpinene (Amiri., 2012; Fachini *et al.*, 2012). Roby *et*

al., (2013) reported that methanolic extracts of thyme are sources of phenolic acids, flavonols and flavanones. Natural foods which are rich in flavonoids helps protect from lung and cancers (Sharangi and Guha., 2013). Eqbal and Abdullah., (2017) confirmed the strong antimicrobial properties of thymol oil which found in thymus essential oil.

According to Ministry of Agriculture Statistics, 2018; the average cultivated area of thymus in the open field was 276 fadden, with a yield quantity of 552 tons. Thymus plant height ranges between 15 to 30 cm and 40 cm wide (Fernanda *et al.*, 2012).

Ascorbic acid (Vitamin C) is essential for plant growth, which is characterized by its antioxidant activity, enzymatic capabilities and promoting power on plant growth. Ascorbic acid is essential in plant photosynthetic activities, cell expansion and resistance to environmental stress, electron transport, plant flowering and senescence,

cell death and fighting pathogens (Blokhina *et al.*, 2003; Gomez and Lajolo, 2008). Many authors confirmed the basic role of ascorbic acid in timing of flowering and senescence (Ahlam and Mustaf, 2019; Carr and Lykkesfeldt., 2020). The foliar application of mixtures (garlic extract, ascorbic acid and nicotinamide) significantly increased plant physical and chemical parameters as reported by Mohamed *et al.*, (2020) on Faba bean.

Amino acids are the building blocks of proteins. Amino acids are of great importance in plant growth which attribute by a large extend in yield enhancement and plant overall development. Naturally extracted amino acids are an important source of nitrogen that can be directly absorbed by plants, as well as it can be considered as major transport forms of organic nitrogen in plants, which have been identified in different plant species (Guangzhe Yang *et al.*, 2020). Foliar application of amino acids was reported to significantly increase yield and plant growth attributes (Khatab *et al.*, 2012; Wassel *et al.*, 2015 and Kamal, 2017).

Active dry yeast as a biofertilizers is reported to has many promoting effects on plant growth. This is attributed to its richness of cytokinin which promotes cell division and synthesis of amino acids and vitamins in plant fruits (Ahmed *et al.*, 2011). In addition to its ability to release CO₂ into soil, which improves soil characteristics and increases plant photosynthetic activity and pigments formation. Many authors have reported the stimulant activities of yeast, of them are Aly *et al.*, (2007) on coriander, Hemdan, (2008) on anise, Dahab *et al.*, (2010) on marjoram and Kenawy, (2010) on Hibiscus sabdariffa, L plants and Matter and. El Sayed (2015) on caraway plants. Active dry yeast foliar application enhanced thyme plants growth and essential oil yield quantity and quality (Heikal, 2005).

Soilless culture has been found to has favorable effects on plant growth and yield quantity and quality (Putra and Yuliando., 2015), off-season crop production (Montagne *et al.*, 2015), introduce new crops which cannot grow on the normal soil (Al-Karaki. and Othman 2009). It improves plant soil interactions and effect on the nutrient availability in the plant root zone which may increase the yield of crops (Montagne *et al.*, 2015). Soilless culture was considered a key alternative in the case of soil borne diseases, which reduces the use of methyl bromide that is used in soil sterilization and led to more environmentally friendly agriculture system (D'Imperio *et al.*, 2018). The type of the used substrate media may significantly affect plant grown (Alsmairat *et al.*, 2018) where media with low bulk density and high-

water holding capacity facilitate plants aeration and root penetration which enhance nutrient availability and absorption (Deepagoda *et al.*, 2013). It has been reported that the substrate physical and chemical properties effect the production quality and quantity such as yield, flower size and number, fruit sugar and phenolic compounds (Al-Ajmi *et al.*, 2009; Schwarz *et al.*, 2009; Al-Ajlouni *et al.*, 2017).

The objective of this study was to determine the growth, yield and chemical composition of *Thymus vulgaris* plant as influenced by the application of different sources of growth regulators in order to improve the yield of herb and essential oil content.

Materials and methods

This experiment was conducted in the Central Laboratory for Agricultural Climate Research Centre (CLAC), Dokki, Giza, Egypt, under unheated greenhouse conditions for two successive seasons of the years 2019 and 2020.

Plant material

At the first week of March, the seedlings were transplanted in plastic pots (5 liters in volume) filled with well mixed media of perlite: sand (1:1 v/v). It was placed on terraces in the greenhouse with drainage capabilities and irrigated by balanced nutrient solution until they formed true leaves. Three weeks later after transplanting, the plants were sprayed with aqueous solution of different biostimulators (vitamin, amino acid (aspartic acids) and dry yeast). The plants were sprayed three times during each cut, the first spray application was occurred after new leaves, the second after one week from the first and the third one after one week from the second one. This sequence was implemented during the second cut and for the second seasons.

Experimental design

Experiment was carried out on a completely randomized design, with 36 blocks arranged in a 1 × 9 × 3 factorial schemes (one cultivar, nine concentrations and three replicated).

Experimental treatments

1. Control (Foliar application with water)
2. Ascorbic acid (1 g/L)
3. Ascorbic Acid (2 g/L)
4. Yeast (2 g/L)
5. Yeast (4 g/L)
6. Amino acid (0.5 g/L): (Aspartic acid 500 ppm/L)
7. Amino acid (1 g/L): (Aspartic acid 1000 ppm/L)

Table a: The chemical composition of nutrient solution used.

Nutrient solution	Macronutrients (ppm)					Micronutrients (ppm)					
	N	P	K	Ca	Mg	Fe	Mn	Zn	B	Cu	Mo
Chemical nutrient solution	200	45	300	180	60	3.0	0.8	0.4	0.5	0.25	0.02

8. Mix1: Foliar application with vitamin at 1 g/l, amino acid (aspartic acids) at 500 ppm and dry yeast at 2 g/l

9. Mix2: Foliar application with vitamin at 2 g/l, amino acid (aspartic acids) at 1000 ppm and dry yeast at 4g/l

The used nutrient solution was adapted from Cooper solution (Cooper, 1979) depending on the analysis of the local water (El-Be hairy, 1994) as shown in table a. The desired initial concentration of the nutrient solution was maintained by suitable dilution of the stock solutions with tap water. Electrical conductivity (EC) of the nutrient solution was maintained between (2–2.2) m.mhos⁻¹ and pH maintained between (6–6.5).

Data sampling and analysis

Sampling: two cuts were harvested, mid- July (First cut) and first week of October (second cut). The plant herbage was harvested by cutting 5 cm above the soil surface and plant growth parameter for the two cuts were recorded as:

Vegetative growth characteristics

- Numbers of branches per plant.
- Plant height (cm).
- Fresh weight (g/herb).
- Dry weight (%): was determined according to A.O.A.C. (1990).

Determination of chemical composition of thymus herbage

Determination of moisture content

The moisture content was calculated according to A.O.A.C. (1990).

Pigments determination

Chlorophyll a, b and carotene were determined according to A.O.A.C. (1990) and Wettstein (1957).

Total carbohydrates

Total carbohydrates were determined according to A.O.A.C. (1990).

Phenolic compounds (mg /100g FW) as catichol

The colorimetric method of Folin-Denis as described by Shahidi and Nacz, (1995).

Essential oil (%)

Quantitative determination of essential oil obtained from different treatments was achieved by hydro-

distillation. The distillation of 100 g fresh herb was continued for 2.5 to 3 hours after water boiling till no further increase in the oil was observed. The oil was permitted to stand undisturbed and the amount of oil obtained from plant materials was calculated.

Oil (%) = (observed volume of oil (ml)/weight of sample (g)) × 100.

Essential oil yield

Essential oil (ml/plant) was calculated by multiplying the average fresh weight of plant by the average oil percentage.

Oil yield/plant = plant fresh weigh (g) × oil %.

Mineral contents (N, P, & K) and Crude protein

Total nitrogen was determined according to kejldahl method as A.O.A.C., (1990). Total phosphorus was determined colorimetrically by ascorbic acid reductant method according to Murphy and Riley, (1962) as modified by Watanabe and Olsen, (1965) and total potassium was determined by using flame photometer according to A.O.A.C, (1990). Crude protein is calculated as mineral nitrogen multiplied by the protein factor, which is 6.25.

GLC analysis of the essential oil

The essential oil constituents were identified by Gas Liquid Chromatography (GLC) according to Radwan (1978).

Economic evaluation

Total cost is the sum of total fixed cost and total variable cost (L.E/m²), where total fixed cost is the sum of the cost of (substrate (sand + perlite) + plastic pots + nutrient solution + irrigation + power) and total variable cost is the sum of the cost of (seedlings + biostimulators + labours). Total revenue was calculated as the sum of the herb fresh weight for the two seasons multiplied by the number of plants in one square meter multiplied by the average market price (L.E./kg). The following table b illustrates the market price of the experiment constituents by Egyptian pound as recorded during the years of 2019 and 2020.

Statistical analysis

The statistical significance of observed differences among treatment means was evaluated by analysis of variance (ANOVA). Statistical analysis was made using

Table b: Price of the experiment constituents (L.E.).

Item	Cost (L.E.)	Item	Cost (L.E.)
Pot	2.5	Seedling	1
Sand (L)	0.30	Nutrient Solution (L)	12
Perlite (L)	1.80	Power (m ²)	10
Irrigation system (m ²)	54	Irrigation (m ²)	10
Tank	50	Ascorbic Acid (1 g)	1
N. plant/m ²	16	Yeast (1 g)	0.37
Thymus Market prices kg/FW	100	Amino acid (1 g)	5

- Net return = Revenue (L.E/m²)- Total cost (L.E/m²)
- The revenue to total Cost (B/C) ratio was calculated to represent the profit percentage.

the “agricolae” package in R software program according to (Duncan, 1955, R, 2017).

Results

The present study has put stress on some biochemical constituents to obtain good idea about the suitable condition for cultivating thymus plant. The bio-stimulatory influence of amino acids (AA), vitamins (Vits), yeast and mixtures combination of them (Mix1 and Mix2) on the different parameters of thymus grown under soilless culture conditions have been investigated herein after.

- Morphological characteristics: plant height (cm), No. of branches/plant, Fresh weight (g/plant), essential

oil % and essential oil (g/herp).

- Chemical characteristics: dry weight%, moisture %, chlorophyll a, b and carotenes (mg/g FW), total pigment, carbohydrate %, total phenols (mg/100 g FW), N, P and K (%) and crud protein.

Effect of different growth promoters on vegetative growth

All treatments have exerted a significant positive influence on plant height, No. of branches/plant, plant fresh and dry weight and moisture content in comparison with the control treatment, which showed the lowest values as

illustrated in table 1 and Fig. 1. Most of these differences




Fig. 1: Productivity (kg/m²) of thymus herb for the two seasons.

Table 1: Effect of different growth promoters on vegetative growth on Thymus Under Soilless Culture Conditions.

Harvest	height (cm)		N. of branches/plant		fresh weight (g/herb)		% Dry weight		% moisture	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
Treatment	First season									
Control	57.67 ^G	61.67 ^G	5.33 ^G	15.00 ^D	88.04 ^H	215.06 ^H	20.99 ^E	21.29 ^F	76.13 ^E	77.57 ^F
Ascorbic acid (1 g/L)	69.61 ^{DE}	74.33 ^{DE}	7.00 ^{DE}	17.00 ^{BC}	108.00 ^{EF}	246.2 ^{EF}	23.42 ^{BCDE}	24.66 ^{CDE}	78.31 ^{BCD}	80.43 ^D
Ascorbic Acid (2 g/L)	74.95 ^{BC}	78.78 ^{BC}	8.00 ^{BC}	18.00 ^{AB}	126.17 ^C	274.75 ^{BC}	25.66 ^B	27.41 ^{ABC}	79.66 ^B	82.29 ^{BC}
Yeast (2 g/L)	63.39 ^F	67.78 ^F	6.00 ^{FG}	16.00 ^{CD}	95.17 ^{GH}	225.5 ^{GH}	21.88 ^{DE}	23.16 ^{EF}	77.00 ^{DE}	78.04 ^{EF}
Yeast (4 g/L)	73.89 ^{BC}	75.78 ^{CD}	8.00 ^{BC}	18.00 ^{AB}	120.41 ^{CD}	265.77 ^{CD}	24.63 ^{BC}	26.19 ^{BCD}	79.54 ^B	82.03 ^{BC}
Amino acid (0.5 g/L)	67.33 ^{EF}	71.33 ^{EF}	6.33 ^{EF}	16.22 ^C	100.25 ^{FG}	233.92 ^{FG}	22.41 ^{CDE}	23.78 ^{DEF}	77.64 ^{CD}	79.12 ^E
Amino acid (1 g/L)	71.56 ^{CD}	74.78 ^{CDE}	7.33 ^{CD}	17.00 ^{BC}	112.70 ^{DE}	258.17 ^{DE}	24.21 ^{BCD}	25.60 ^{BCDE}	78.52 ^{BC}	81.47 ^{CD}
Mix 1	76.97 ^B	81.94 ^B	8.89 ^B	18.00 ^{AB}	139.83 ^B	282.08 ^B	26.05 ^B	27.77 ^{AB}	81.36 ^A	82.86 ^{AB}
Mix 2	83.08 ^A	86.90 ^A	10.00 ^A	18.67 ^A	162.75 ^A	302.5 ^A	30.19 ^A	30.12 ^A	81.48 ^A	83.51 ^A
Treatment	Second season									
Control	66.60 ^F	66.33 ^G	5.00 ^E	14.33 ^D	96.26 ^H	226.8 ^G	21.92 ^E	20.89 ^F	77.21 ^F	77.37 ^D
Ascorbic acid (1 g/L)	71.96 ^E	75.47 ^{EF}	6.00 ^{DE}	16.00 ^{BC}	122.18 ^{EF}	247.41 ^{DE}	24.26 ^{CDE}	23.47 ^{DE}	80.13 ^{CDE}	79.22 ^{BC}
Ascorbic Acid (2 g/L)	80.22 ^{BC}	83.58 ^{BC}	7.67 ^{BC}	17.33 ^A	150.98 ^{BC}	273.69 ^B	26.04 ^{BC}	26.23 ^{BC}	82.75 ^A	81.17 ^A
Yeast (2 g/L)	69.45 ^{EF}	71.62 ^F	5.33 ^E	15.00 ^{CD}	104.72 ^{GH}	235.56 ^{FG}	22.57 ^E	22.07 ^{EF}	78.43 ^{EF}	78.17 ^{CD}
Yeast (4 g/L)	78.45 ^{CD}	81.20 ^{CD}	7.00 ^{CD}	17.00 ^{AB}	138.09 ^{CD}	262.02 ^C	25.41 ^{BCD}	25.00 ^{CD}	81.57 ^{ABC}	80.36 ^{AB}
Amino acid (0.5 g/L)	70.84 ^E	73.94 ^{EF}	6.00 ^{DE}	16.00 ^{BC}	113.50 ^{FG}	240 ^{EF}	23.33 ^{DE}	22.90 ^{DEF}	79.44 ^{DE}	78.30 ^{CD}
Amino acid (1 g/L)	75.37 ^D	77.57 ^{DE}	7.00 ^{CD}	16.89 ^{AB}	127.98 ^{DE}	252.87 ^{CD}	24.94 ^{BCD}	24.42 ^{CD}	80.51 ^{BCD}	79.38 ^{BC}
Mix 1	81.86 ^B	86.69 ^{AB}	8.22 ^B	18.00 ^A	159.31 ^{AB}	283.16 ^{AB}	26.79 ^B	27.38 ^{AB}	83.18 ^A	81.58 ^A
Mix 2	86.08 ^A	88.87 ^A	9.67 ^A	18.00 ^A	171.43 ^A	292.91 ^A	30.82 ^A	29.33 ^A	82.02 ^{AB}	81.51 ^A

did not reach level of significance among each other. The highest values of plant height, No. of branches/plant and fresh weight have been attained in case of spraying with the highest concentrations of Vit. C, amino acids and yeast mixtures (Mix2). While, mixture of low concentration (Mix1) came at the second order in enhancing the vegetative growth. Plant height reached approximately about 1.42 and 1.32 times the corresponding control in the first and second seasons respectively. While, No. of branches/plant were approximately (1.56 and 1.59) times the control treatment in the first and the second seasons respectively. Foliar application of Mix2 led to an increase in plant fresh weight by 1.63 and 1.53 g/herb for the two seasons respectively. Fig. 1 represents the plant productivity response to the different treatments where the mixture treatments were the highest during the two studied seasons.

Concerning dry weight (%) and moisture content (%) of thymus plants, data illustrated in table 1 reveals that, the highest values recorded were due to the application of Mix treatments with a significant difference compared to all other treatments including the control one. This was followed by the application of ascorbic acid which was significantly different from the control treatment but, did not reach the level of significance compared to the other treatments.

Effect of different growth promoters on Essential oil %, essential oil (g/herp), total carbohydrates and total phenol

It is well known that the economic value of thymus plant is attributed to its essential oil content therefore we have put stress on the positive variations that may be resulted due to cultivating thymus plant in perlite: sand (1:1 v/v) soilless culture and the regulatory effect of growth promoters (AA, vit. C, yeast and their mixtures). Thus, the evaluation and comparison among all treatments were based on essential oil percentage together with its yield (ml oil/plant fresh weight). Data obtained are listed in table 2 for the first and second cut during the two studied seasons. Data reveals that thymus plants responded positively to the application of all biostimulators however this increase in the essential oil content did not reach the level of significance compared with the control treatment. The highest significant increase in the oil content was a result of the mix application followed by the ascorbic acid treatment. This holds true for the two cuts during the two seasons.

The higher concentration mixture of the applied biostimulators (Mix2) gave the highest values during the two cuts and the two successive seasons (0.5 and 0.58

%). While the other mixture (Mix1) came in the second order (0.37 and 0.47 %). The difference between Mix2 and Mix1 in essential oil was significantly different with each other and with the control treatment. The lowest essential oil percentage and essential oil (ml/herb fresh weight) was recorded in case of the control treatment (0.19 – 0.26%). This holds true for the two successive cuts during the two seasons. The marvelous results have been achieved by foliage spraying with Mix2, which produced highest significant essential oil % and essential oil (g/herp), which reached 2.63 – 2.2 essential oil% and 4.8 – 4.4 essential oil (g/herp) times than the corresponding control respectively, in the first and second season.

Data presented in table 2 shows the responses of total carbohydrates (%) and total phenols (mg/100g FW) responses due to the foliar spray of the investigated biostimulators for the two cuts during both seasons. Generally, all treatments increased the total carbohydrates and phenols in thymus plants. However, this increase did not reach the level of significance in some treatments such as yeast and amino acids treatments regardless the concentration compared to the control treatment. The Mix treatments have excreted a significant increase of both carbohydrates and phenols of thymus plants compared to all other treatments including the control one. But the differences between the two treatments was not almost significantly from cut to another or season to season with the highest value obtained in the case of Mix2 treatment. This holds true for the two cuts during the two studied seasons.

Effect of different growth promoters on GC profile of thymus essential oil

The GC profile of thymus essential oil reveals the presence of eleven identified compounds of them, ten were existed in relatively minute proportions (minor components). While thymol oil was the major found oil as illustrated in tables 3a and 3b for the two seasons respectively.

The main features that may be characterized the essential oil investigated in the present study which extracted from thymus plants grown in perlite: sand (1:1 v/v) soilless culture, are:

- The superiority of thymol, which varied from one cut to another. The highest value was attained by Mix2 (53.42 – 53.22 %) compared with all other treatments. Whilst the lowest value (44.49 – 46.102%) recorded in the control in the first and second cuts respectively.

- ρ -cymene was the second major constituent in all treatments and showed variable fluctuations due to growth promoters application. The control recorded (9.85 – 9.84

Table 2: Effect of different growth promoters on Essential oil, Total carbohydrates and Total phenol on Thymus Under Soilless Culture Conditions.

Harvest	Essential Oil %		Oil (g/herp)		%Total Carbohydrates		Toal Phenols	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
Treatment	Frist season							
Control	0.19 F	0.26 F	0.17 E	0.36 E	5.50 G	5.56 F	2.37 E	1.91 E
Ascorbic acid (1 g/L)	0.24 DEF	0.37 CD	0.26 CDE	0.65 CDE	6.07 EF	6.24 E	2.76 DE	2.25 DE
Ascorbic Acid (2 g/L)	0.33 BC	0.42 BC	0.41 BC	1.13 BC	7.03 C	7.49 BC	3.75 BC	3.13 BC
Yeast (2 g/L)	0.21 EF	0.3E F	0.20 E	0.45 DE	5.66 FG	5.90 EF	2.49 DE	1.99 E
Yeast (4 g/L)	0.31 BCD	0.4 C	0.37 BCD	0.98 BCD	6.75 CD	7.03 CD	3.27 CD	2.71 CD
Amino acid (0.5 g/L)	0.23 DEF	0.32 DE	0.23 DE	0.54 DE	5.76 EFG	5.99 EF	2.59 DE	2.08 DE
Amino acid (1 g/L)	0.28 CDE	0.38 CD	0.32 CDE	0.82 CDE	6.32 DE	6.44 DE	2.95 DE	2.51 CDE
Mix 1	0.37 B	0.47 B	0.51 B	1.45 B	7.70 B	7.91 AB	4.20 B	3.55 B
Mix 2	0.5 A	0.58 A	0.82 A	2.5 A	8.37 A	8.40 A	5.60 A	4.80 A
	Second season							
Control	0.21 F	0.25 F	0.21 F	0.47 F	4.35 F	5.56 F	2.84 F	2.28 F
Ascorbic acid (1 g/L)	0.27 CDEF	0.31 CDE	0.33 DEF	0.81 DEF	5.06 EF	6.24 E	3.75 E	3.10 DE
Ascorbic Acid (2 g/L)	0.34 BC	0.37 BC	0.51 BC	1.4 BC	6.81 BC	7.49 BC	4.78 BC	3.75 BC
Yeast (2 g/L)	0.23 EF	0.27 EF	0.24 EF	0.58 EF	4.58 F	5.90 EF	3.35 EF	2.74 EF
Yeast (4 g/L)	0.32 BCD	0.35 BCD	0.45 CD	1.17 CD	6.18 CD	7.03 CD	4.37 CD	3.54 BCD
Amino acid (0.5 g/L)	0.25 DEF	0.30 DEF	0.29 DEF	0.69 DEF	4.81 F	5.98 EF	3.58 E	2.97 DE
Amino acid (1 g/L)	0.30 CDE	0.34 CD	0.39 CDE	0.98 CDE	5.79 DE	6.44 DE	3.92 DE	3.23 CDE
Mix 1	0.39 B	0.40 B	0.62 B	1.76 B	7.18 B	7.91 AB	5.07 B	4.15 B
Mix 2	0.50 A	0.58 A	0.87 A	2.54 A	8.16 A	8.40 A	5.91 A	4.84 A

%) in first and second cut respectively.

· Carvacrol low values compared with γ -terpinene and α -Pinene recorded.

Effect of different growth promoters on N, P and K Contents (% of dry matter) and crud protein

Data illustrated in table 4 show that the effect of

Table 3a: Effect of different growth promoters on GC profile of Thymus Essential oil First Season.

Treatment	com- pone nts %	α - Pin ene	Sab ine ne	myr cene	ρ - cym ene	α - terpi- nnene	Bor neol	γ - terp inene	thy mol	car vac rol	und eca nol	α - Hun ulene
Control	1 st	1.47	8.16	3.036	9.86	3.39	5.486	7.75	44.49	6.76	3.702	4.49
	2 nd	1.16	8.3	5.61	9.84	6.06	5.25	7.17	46.102	6.702	1.341	2.25
Ascorbic acid (1 g/L)	1 st	0.86	6.48	2.24	10.82	0.65	4.19	6.02	47.91	8.83	3.73	4.14
	2 nd	1.47	6.16	3.03	9.86	1.39	5.48	7.75	48.97	7.76	5.49	3.7
Ascorbic Acid (2 g/L)	1 st	0.89	6.56	2.07	9.82	2.55	4.63	7.87	48.22	8.4	4.04	3.47
	2 nd	1.39	6.85	2.51	9.87	1.809	5.47	7.98	49.66	6.09	2.051	3.67
Yeast (2 g/L)	1 st	0.59	6.26	5.34	8.83	5.57	4.68	7.38	49.42	6.89	2.25	0.745
	2 nd	0.811	6.67	2.49	8.18	5.41	6.35	8.09	48.018	5.83	2.11	2.02
Yeast (4 g/L)	1 st	0.76	5.88	3.32	8.29	2.32	3.41	8.9	52.04	7.24	0.63	1.35
	2 nd	1.4	8.9	3.06	6.29	2.37	6.63	7.3	53.29	6.29	1.82	0.99
Amino acid (0.5 g/L)	1 st	0.53	1.912	8.01	9.92	4.78	0.007	7.63	50.3	8.79	0.68	1.24
	2 nd	0.48	1.66	10.042	10.29	3.322	0.021	8.23	50.21	6.15	0.63	0.26
Amino acid (1 g/L)	1 st	0.74	5.74	1.77	9.4	4.68	6.67	7.38	50.91	6.22	0.61	1.45
	2 nd	1.61	8.45	2.7	8.14	7.29	4.59	7.23	51.19	5.27	1.132	1.15
Mix 1	1 st	0.86	6.48	2.24	10.82	0.65	4.19	6.024	48.91	8.83	3.73	4.17
	2 nd	0.85	5.1	4.47	9.44	5.68	6.04	9.36	50.08	4.89	1.36	1.36
Mix 2	1 st	0.75	5.3	6.9	9.3	6.7	4.9	7.14	53.41	5.35	0.14	1.21
	2 nd	0.67	5.98	6.47	8.86	6.54	5.07	8.61	53.22	5.39	1.42	1.42

Table 3b: Effect of different growth promoters on GC profile of *Thymus* Essential oil Second Season.

Treatment	com- pone nts%	α - Pin ene	Sab ine ne	myr cene	ρ - cym ene	α - terpi- nnene	Bor neol	γ - terp inene	thy mol	car vac rol	und eca nol	α - Hun ulene
Control	1 st	1.55	2.93	8.03	9.94	4.80	7.20	7.65	44.32	8.81	0.70	1.26
	2 nd	1.52	2.90	8.00	9.91	4.77	7.17	7.62	42.40	8.78	0.67	1.23
Ascorbic acid (1 g/L)	1 st	1.17	6.58	2.50	9.82	2.68	5.10	7.03	52.31	7.45	3.36	3.05
	2 nd	1.17	6.41	2.59	10.17	1.57	4.92	6.93	50.20	8.01	4.19	3.63
Ascorbic Acid (2 g/L)	1 st	1.15	6.61	2.39	9.95	1.98	5.01	7.59	48.87	7.50	3.43	3.59
	2 nd	1.14	6.67	2.32	9.88	2.11	5.04	7.81	51.23	7.33	3.17	3.58
Yeast (2 g/L)	1 st	0.53	1.91	8.01	9.92	4.78	6.76	7.63	50.02	8.79	0.68	1.24
	2 nd	0.81	2.67	8.20	8.18	5.41	6.35	7.03	51.34	5.83	2.11	2.02
Yeast (4 g/L)	1 st	1.29	1.15	8.77	9.02	5.54	7.52	8.39	50.78	6.55	1.44	2.00
	2 nd	1.99	1.87	6.56	9.13	4.60	4.34	8.49	52.82	8.56	1.27	2.14
Amino acid (0.5 g/L)	1 st	0.75	3.66	6.77	9.01	4.18	1.66	7.55	51.05	8.17	0.97	1.18
	2 nd	0.61	2.66	8.41	9.65	3.75	0.84	7.89	50.63	7.16	0.80	0.72
Amino acid (1 g/L)	1 st	0.68	4.20	5.09	9.53	4.21	3.76	7.63	52.03	6.69	0.70	1.08
	2 nd	1.18	7.10	2.24	8.77	5.99	5.63	7.31	52.05	5.75	0.87	1.30
Mix 1	1 st	0.95	6.08	3.01	10.05	2.81	5.09	7.73	50.23	7.02	2.30	2.58
	2 nd	0.89	5.89	3.24	10.10	3.05	5.11	7.71	51.04	6.91	2.01	2.25
Mix 2	1 st	0.47	3.76	4.46	6.05	4.41	3.32	5.25	53.45	6.58	3.63	3.33
	2 nd	0.63	5.01	5.94	8.07	5.88	4.43	6.25	52.87	7.73	2.89	0.54

different treatments on thymus contents of N, curd protein, P and K %. Data reveals that the Mix and ascorbic acid treatments excreted a significant increase on the herb contents of N, P, K and crud protein in the first and second cuts during the two studied seasons. The other treatments increased the NPK and crud protein compared with the control treatment however this increase did not reach the level of significance during the two seasons.

The NPK and crud protein content of thymus responded differently to the concentration of the added bioregulators where Mix2 led to a significant increase compared with Mix1. Also, the higher concentration of yeast (4 g/L) led to a significant increase compared with the lower one (2 g/L). This holds true for NPK and crud protein contents for both cuts during both seasons.

Effect of different growth promoters on Chlorophyll a, chlorophyll b, carotene and total pigments content

All photosynthetic pigments increased due to the application of growth promoters compared with the control treatment however this increase did not reach the level of significance. Data in table 5 illustrate the response of Chl. a, b, carotene and total pigments to the application of different treatments. Data reveals that the Mix and ascorbic acid treatments excreted a significant increase on the herb contents of all photosynthetic pigments in the first and second cuts during the two studied seasons. The other treatments increased the

Chlorophyll a, chlorophyll b, carotene and total Pigments content compared with the control treatment however this increase did not reach the level of significance during the two seasons.

The Chlorophyll a, chlorophyll b, carotene and total Pigments content of thymus responded differently to the concentration of the added bioregulators where Mix2 led to a significant increase compared with Mix1. Also, the higher concentration of yeast (4 g/L) led to a significant increase compared with the lower one (2 g/L). This holds true for the Chlorophyll a, chlorophyll b, carotene and total Pigments content for both cuts during both seasons.

Economic Evaluation


The economic feasibility of this experiment was carried out and the net return and benefit cost ratio (B/C) of the field experiment for thymus fresh leaves are shown in table 6 and Fig. 2. The results showed that the ascorbic acid (2 g/l) came in the first order in terms of (B/C) where it scored 1.8 B/C, Yeast (4 g/L) came at second where it scored 1.73 in terms of B/C ratio.

Discussion

In this experiment, results showed the positive influence of yeast, amino acids, vitamins and their mixtures on thymus herbs quality and quantity. Many authors agree with these findings such as Abou EL-Yazied and Mady (2012) on bean plants and Mahmoud, *et al.*, (2016) on lupine (*Lupinus termis* L.) plants, where they

Table 4: Effect of different growth promoters on Mineral Content on Thymus Under Soilless Culture Conditions.

Harvest	% Nitrogen		% crud protein		% Phosphorus		% potassium	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
Treatment	Frist season							
Control	1.44F	1.41F	9.02F	8.83F	0.41F	0.50F	1.13F	0.94G
Ascorbic acid (1 g/L)	1.72DE	1.83CD	10.73DE	11.42CD	0.6CDE	0.59DE	1.42CD	1.18DE
Ascorbic Acid (2 g/L)	1.94BC	2.04B	12.10BC	12.75B	0.69B	0.65BC	1.55BC	1.3C
Yeast (2 g/L)	1.49F	1.60EF	9.29F	10.02EF	0.52E	0.55E	1.22EF	1.07F
Yeast (4 g/L)	1.83CD	1.99BC	11.42CD	12.46BC	0.67BC	0.64BCD	1.51C	1.25CD
Amino acid (0.5 g/L)	1.65E	1.70DE	10.29E	10.60DE	0.57DE	0.56E	1.3DE	1.12EF
Amino acid (1 g/L)	1.77DE	1.94BC	11.08DE	12.10BC	0.65BCD	0.62CD	1.46C	1.22CDE
Mix 1	2.05B	2.11B	12.81B	13.19B	0.71B	0.68B	1.65AB	1.41B
Mix 2	2.30A	2.34A	14.35A	14.60A	0.81A	0.78A	1.78A	1.54A
	Second season							
Control	1.29 F	1.16 G	8.09 F	7.24 G	0.35 F	0.47 F	1.12 F	1.06 F
Ascorbic acid (1 g/L)	1.51 DE	1.39 DEF	9.41 DE	8.67 DEF	0.44 CDE	0.56 DE	1.57 CD	1.28 DE
Ascorbic Acid (2 g/L)	1.79 C	1.56 BC	11.17 C	9.76 BC	0.49 BC	0.60 BC	1.77 B	1.59 BC
Yeast (2 g/L)	1.42 EF	1.30 F	8.87 EF	8.11 F	0.38 EF	0.50 F	1.38 E	1.21 EF
Yeast (4 g/L)	1.72 C	1.50 CD	10.74C	9.38 CD	0.47BCD	0.59BCD	1.72 BC	1.52 BC
Amino acid (0.5 g/L)	1.47 EF	1.32 EF	9.18 EF	8.25 EF	0.41 DE	0.55 E	1.46 DE	1.23 EF
Amino acid (1 g/L)	1.66 CD	1.44CDE	10.36CD	9.00CDE	0.46BCD	0.58CDE	1.67 BC	1.42 CD
Mix 1	2.00 B	1.68 B	12.48B	10.50B	0.52 B	0.62 AB	1.82 AB	1.66 B
Mix 2	2.29 A	1.85 A	14.29A	11.54A	0.61 A	0.66 A	1.94 A	1.86 A

**Fig. 2:** Relationship between operating costs and net revenue of Thymus plants under soilless culture conditions.

confirmed the significant increase in plants vegetative growth when sprayed with yeast. The superiority of Mix2 treatment may be attributed due to its richness of amino acids which are considered as the building blocks in cell biosynthesis, improve various biochemical processes and facilitate nutrients availability (Dawood and Sadak 2007). Moreover, this mixture is rich in its content of macro and micro nutrients due to existence of high concentrations of yeast, vitamins and amino acids which may play an important role in improving plant vegetative growth and increases plant resistance to diseases (Bevilacqua, *et al.*, 2008). The stimulant role of yeast as a growth promoter was attributed to the role it may plays in cell division and photosynthetic activities such as pigments formation, protein and nucleic acid synthesis. Moreover, yeast has been found to contains protective agents such as amino acids, proteins and vitamins that catalysis plant growth (Wanas, 2006).

The application of Mix treatments were significantly increased plant vegetative parameters with the superiority of Mix2 that led to an average increase of 1.5 times the control treatment of plant fresh weight. Many authors found that foliar application of vitamins significantly increased plant height, No. of branches/plant, fresh, dry weight and essential oil percentage of plants, of them are Fatma *et al.*, (2008 and 2014) on lavender and strawberries plants respectively, Nahed *et al.*, (2010) on *Thuja orientalis* L., Hendawy *et al.*, (2010; 2015) on Thymus and *Mentha piperita* plants respectively. Azza Ezz El-Din and Hendawy, (2010) confirmed the superiority of the higher concentration of ascorbic acid on *Borago officinalis* plants vegetative growth such as plant height, number of flowers, number of branches and seed weight. While Shafeek *et al.*, (2015) and Ahmed and Farm (2015) reported the significant increase vegetative growth of garlic plants and Fawzy *et al.*, (2012) on onion plant when sprayed with EM, yeast and amino acids respectively.

All treatments increased thymus essential oil with the superiority of Mix treatments, where Mix2 doubled thymes essential oil percentage compared to the control treatment. These results are similar with many studies were the positive effect of plant growth bio-stimulators on essential oil percentage and oil (ml/herb) were reported (Aly *et al.*, 2007; Hemdan, 2008; Dahab *et al.*, 2010; Kenawy, 2010). All treatments increased the carbohydrate

Table 5: Effect of different growth promoters on pigment content on *Thymus* Under Soilless Culture Conditions.

Harvest	Ch. A (mg /100 FW)		Ch. B (mg /100 FW)		Carotene (mg/100 FW)		Total pigment (mg/100 FW)	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
Treatment	Frist season							
Control	0.11F	0.13F	0.07F	0.05E	0.05D	0.04F	0.23G	0.23E
Ascorbic acid (1 g/L)	0.18CD	0.16DE	0.08DE	0.08CD	0.06D	0.07CD	0.32DE	0.31D
Ascorbic Acid (2 g/L)	0.20BC	0.21B	0.10BC	0.11B	0.08BC	0.09B	0.39BC	0.41B
Yeast (2 g/L)	0.14EF	0.13F	0.08EF	0.07D	0.05D	0.05EF	0.27FG	0.25E
Yeast (4 g/L)	0.20BC	0.19BC	0.10CD	0.09C	0.08BC	0.09B	0.37C	0.37BC
Amino acid (0.5 g/L)	0.16DE	0.15EF	0.08EF	0.08CD	0.06D	0.06DE	0.3EF	0.29D
Amino acid (1 g/L)	0.19CD	0.18CD	0.09CDE	0.09C	0.07C	0.08BC	0.35CD	0.35C
Mix 1	0.22B	0.23A	0.12AB	0.12AB	0.09AB	0.11A	0.43B	0.46A
Mix 2	0.28A	0.24A	0.12A	0.13A	0.09A	0.12A	0.49A	0.5A
	Second season							
Control	0.13 F	0.12 G	0.08 G	0.05 D	0.05 G	0.06 F	0.25 G	0.23 F
Ascorbic acid (1 g/L)	0.16 DE	0.17 DE	0.09 EF	0.11 D	0.06 EF	0.08 DE	0.31 EF	0.36 E
Ascorbic Acid (2 g/L)	0.19 BC	0.21 BC	0.11 BC	0.56 AB	0.08 BC	0.10 B	0.38 BC	0.86 BC
Yeast (2 g/L)	0.14 EF	0.14 FG	0.08 FG	0.07 D	0.05 G	0.07 EF	0.27 G	0.27 EF
Yeast (4 g/L)	0.18 C	0.19 CD	0.10 CD	0.50 B	0.07 CD	0.09 BC	0.35 CD	0.79 C
Amino acid (0.5 g/L)	0.15 EF	0.15 EF	0.08 EFG	0.08 D	0.05 FG	0.07 EF	0.29 FG	0.30 EF
Amino acid (1 g/L)	0.17 CD	0.19 CD	0.09 DE	0.38 C	0.06 DE	0.08 CD	0.33 DE	0.65 D
Mix 1	0.20 B	0.24 B	0.12 AB	0.58 AB	0.08 B	0.11 A	0.41 B	0.93 AB
Mix 2	0.23 A	0.28 A	0.13 A	0.62 A	0.09 A	0.12 A	0.46 A	1.02 A

Table 6: Economic evaluation of field experiment (leaves of *Thymus* crop) Under soilless culture conditions (L.E./m²).

Treatment	Input			Output			
	Fixed Cost (L.E/m ²)	Variable Cost (L.E/m ²)	Total cost (L.E/m ²)	Economic criterion			
				Revenue (L.E/m ²)	Net Revenue (L.E/m ²)	B/C ratio	Treatment Order
Control	254	178	432	1001.86	569.86	1.32	7
Ascorbic acid (1 g/L)	254	198	452	1158.06	706.06	1.56	4
Ascorbic Acid (2 g/L)	254	218	472	1320.94	848.94	1.80	1
Yeast (2 g/L)	254	192.8	446.8	1057.52	610.72	1.37	6
Yeast (4 g/L)	254	207.6	461.6	1258.06	796.46	1.73	2
Amino acid (0.5 g/L)	254	228	482	1100.27	618.27	1.28	8
Amino acid (1 g/L)	254	278	532	1202.75	670.75	1.26	9
Mix1	254	262.8	516.8	1383.01	866.21	1.68	3
Mix2	254	347.6	601.6	1487.34	885.74	1.47	5

contents of thymus plants in comparison with the control treatment. The application of Mix2 treatment led to an average increase of 1.5 times the control carbohydrates. Total carbohydrate contents results are similar to those of El-Sherbeny *et al.*, (2012) on *Brassica rapa*, Ali, *et al.*, (2014) (2017) on turnip and Garlic plants, Hanafy *et al.*, (2017) on *Artemisia abrotanum* and El-Khateeb *et al.*, (2017), Hanafy *et al.*, (2018) on *Majorana* plants where they reported the significant increase of total carbohydrates compared to the control treatment when plants were sprayed with some amino acids.

For GC essential oil profile our results confirmed the

dominance of thymol among all other constituents which is in agreement with Aly *et al.*, (2007), Hemdan (2008), Dahab *et al.*, (2010), Kenawy (2010), Maher *et al.*, (2011), Olga Kosakowska *et al.*, (2020) and Shahad *et al.*, (2020). According to the total phenols content; one can find that; mixture treatments increased the total phenols by an average of 2.25 and 1.8 for Mix2 and Mix1 respectively. Many authors agree with these results and confirm the importance of increasing total phenols which may be reflected as increasing in the antioxidant power (Prasanth Reddy *et al.*, 2014; Kocira., 2019; Gema Nieto., 2020). The results of increasing pigments content

with mixture treatments are in conformity with that recorded by Refaat and Balbaa (2001) on lemongrass, Hassanain *et al.*, (2006) on *Matricaria chamomilla* and Naglaa., (2020) *Helianthus annuus*.

Results of this study, directly indicate economic feasibility of the application of plant growth bio-stimulators, which are extremely important to the farmers (Kocira *et al.*, 2020). In contrast to the physical and chemical response of thymus plant to the used bio-stimulators where the mixture treatments were the highest among all other treatments, ascorbic acid treatment (2g/l) and yeast (4 g/l) were the most profitable treatments. this may be attributed to the difference of the variable cost where the mixture treatments variable cost is high.

Conclusion

From the obtained results it can be concluded that, foliar spraying of mixtures of bio-stimulators enhanced the growth, oil percentage and oil yield of thymus. Based on the trial results, it is possible recommended that treating thymus plants with heigh concentrations mixture of AA + Vit. C + yeast significantly increased the yield, productivity, high oil yield. However, from the economic evaluation due to the high cost of this mixture it was not the highest in terms of net return. While the spray with yeast (4 g/l) came at the first order in the B/C ratio due to the low cost of yeast. Thymus cultivation are economically profitable including the control treatment. The bio-stimulants could result in environmentally safe plants to minimize the hazards of pollution caused by using mineral fertilizers.

References

- A.O.A.C (1990). Official methods of analysis of the association of Official analytical chemist's 15th Ed. Published by the association of official analytical chemists, INC suite 400. 2200 Wilson Boulevard, Arlington, Virginia. 22201 USA.
- Abou El-Yazied A. and M.A. Mady (2012). Effect of boron and yeast extract foliar application on growth, pod setting and both green and seed yield of broad bean (*Vicia faba* L.). *J. Am. Sci.*, **8**: 517-534.
- Abou El-Yazied A. and M.A. Mady (2012). Effect of boron and yeast extract foliar application on growth, pod setting and both green and seed yield of broad bean (*Vicia faba* L.). *J. Am. Sci.*, **8**: 517-534.
- Ahlam, A.H. and K. Mustaf (2019). Responses of two varieties of *Cucurbita pepo* L. Planted Inside the Plastic Houses to foliar spraying of ascorbic acid. *Plant Archives*, **19**(1): 1768-1772.
- Ahmed, M.A., M.S. Shalaby, M.S. Sadak, K.M. Gamal El-Din, Y.R. Abdel-Baky and M.A. Khater (2016). Physiological role of antioxidant in improving growth and productivity of chickpea (*Cicer arietinum* L.) grown under newly reclaimed sandy soil. *Res. J. Pharm. Biol. Chem. Sci.*, **7**(6): 399-409.
- Ahmed, M.E.M. and E. Farm (2015). Response of garlic plants (*Allium sativum* L.) to foliar application of some biostimulants. *Egypt J. Hort.*, **42**: 613-625.
- Ahmed, A.A., M.M.H. Abd EL-Baky, M.F. Zaki and Faten S Abd EL-Aal (2011). Effect of foliar application of active yeast extract and zinc on growth, yield and quality of potato plant. *J. of App. Sci., Res.*, **7**(12): 2479-2488.
- Ahmed, A.F., F.A. Attia, Z. Liu, C. Li, J. Wei and W. Kang (2019). Antioxidant activity and total phenolic content of essential oils and extracts of sweet basil (*Ocimum basilicum* L.) plants. *Food Sci. Hum. Wellness*, **8**: 299-305.
- Al-Ajlouni, M., J. Ayad and Y. Othman (2017). Particle size of volcanic tuff improves shoot growth and flower quality of Asiatic hybrid lily using soilless culture. *Hort. Technol.*, **27**: 223- 227.
- Al-Ajmi, A., G. Al-Karaki and Y. Othman (2009). Effect of different substrates on fruit yield and quality of cherry tomato grown under recalculating soilless system. *Acta Hort.*, **2**(807): 491-494.
- Alexandre, P. and I.O.G. Ronnel (2007). Chemical composition of *Thymus vulgaris* L. (thyme) essential oil from the Rio de Janeiro State (Brazil).
- Ali, A.H., M.R. Shafeek, A.R. Mahmoud and M. El- Desuki (2014). Effect of various levels of organic fertilizer and humic acid on the growth and roots quality of turnip plants (*Brassica rapa*). *Curr. Sci. Int.*, **3**(1): 7-14.
- Ali, MAM (2017). Effect of some Bio-stimulants on Growth, Yield and Bulb Quality of Garlic Grown in Newly Reclaimed Soil, New Valley-Egypt. *J. Plant Production, Mansoura Univ.*, **8**(12): 1285 - 1294.
- Al-Karaki, G and Y. Othman (2009). Soilless cultivation of some medicinal and aromatic her plants under the conditions of Arabian Gulf region. *Emirates J. Food Agr.*, **21**(2): 64-70.
- Alsmairat, N., M. Al-Ajlouni, J. Ayad, Y. Othman and R. Hilaire (2018). Composition of soilless substrates affect the physiology and fruit quality of two strawberry (*Fragaria × ananassa* Duch.) cultivars. *J. Plant Nutr.*, **41**: 1420-1430.
- Aly, M.K., M.A. Mohamed, F.A. Attia, A.A. El-Sayed and M.H. Abd El-Gawad (2007). Effect of some organic and biofertilization treatments on coriander plants. II- Volatile oil and chemical composition. 1st Inter. Conf Desert. Cultivation, Problem & Solutions. Minia Univ. p.113-125.
- Amiri, H. (2012). Essential oils composition and antioxidant properties of three *Thymus* species. Evidence-Based Complementary and Alternative Medicine, vol., Article ID 728065, 2012.
- Aouam, I., Y. El Atki, A. Taroq, F. El Kamari and I.A. Abellio (2019). Chemical composition, antimicrobial and antioxidant activities of two Moroccan *Thymus* essential oils. *Asian J. Pharm. Clin. Res.*, **12**: 447-451.

- Azza E. EL-Din and S.F. Hendawy (2010). Effect of dry yeast and compost tea on growth and oil content of *Borago officinalis* plants. *Research Journal of Agriculture and Biological Science*, **6(4)**: 424-430.
- Bakry, B.A., F.M. Ibrahim, M.M.S. Abdallah and H.M.S. El-Bassiouny (2016). Effect of banana peel extract or tryptophan on growth, yield and some biochemical aspects of quinoa plants under water deficit. *International Journal Pharm. Tech. Research*, **9(8)**: 276-287.
- Bevilacqua, A., M.R. Corbo, M. Mastromatteo and M. Sinigaglia (2008). Combined effects of pH, yeast extract, carbohydrates and di-ammonium hydrogen citrate on the biomass production and acidifying ability of a probiotic *Lactobacillus plantarum* strain, isolated from table olives, in a batch system. *World J. Microbiol Biotechnol.*, **24**: 1721-1729.
- Blokhina, O., E. Virolainen and K.V. Fagerstedt (2003). Antioxidants, oxidative damage and oxygen deprivation stress. *Ann. Bot.*, **91**: 179-194.
- Carr, A.C. and J. Lykkesfeldt (2020). Discrepancies in global vitamin C recommendations: A review of RDA criteria and underlying health perspectives. *Crit. Rev. Food Sci. Nutr.*, 1-14.
- Cooper, Allen (1979). The ABC of NFT Grower Books. London. p: 181.
- D'Imperio, M., F. Montesano, M. Renna, B. Leoni, D. Buttarò, A. Parente and F. Serio (2018). NaCl stress enhances silicon tissue enrichment of hydroponic "baby leaf" chicory under biofortification process. *Sci. Hort.*, **235**: 258-263.
- Dahab, T.A.M., I.M.A. Harridy and B.A.B. Mansour (2010). Effect of irrigation and anti-transparent treatments on growth, yield and chemical constituents of marjoram plants (*Majorana hortensis* Moench). *Bulletin of Fac. of Agric. Cairo Univ.*, **61(3)**: 274-285.
- Dawood, M.G., Y.R. Abdel-Baky, M.E. El-Awadi and G.S. Bakhom (2019). Enhancement quality and quantity of faba bean plants grown under sandy soil conditions by nicotinamide and/or humic acid application. *Bulletin of the National Research Centre*, **43**: 28.
- Dawood, M.G. and M.Sh. Sadak (2007). Physiological response of canola plants (*Brassica napus* L.) to tryptophan or benzyl adenine. *Lucrari Stiintifice*, **50(9)**: 198-207.
- Deepagoda, T.K., J. Lopez, P. Møldrup, L. De Jonge and M. Tuller (2013). Integral parameters for characterizing water, energy and aeration properties of soilless plant growth media. *J. Hydrol.*, **502**: 120-127.
- Duncan, D.B. (1955). Multiple range and multiple "F" test. *Biometrics*, **11**: 1-42.
- El-Awadi, M.E., G. Mona, Y.R. Dawood and E.A.H. Abdel-Baky (2017). Physiological effect of melatonin, IAA and their precursor on quality and quantity of chickpea plants grown under sandy soil conditions Agricultural, Engineering. International, special Issue 35-44.
- El-Bassiouny H.M.S., A.A. Abd El-Monem, M.S. Sadak and N.M. Badr (2017). Amelioration of the adverse effects of salinity stress by using ascorbic acid in sunflower cultivars. *Bull. NRC*, **41(2)**: 233-249.
- El-Behairy, U. (1994). The effect of levels of phosphorus and zinc in the nutrient solution on macro and micronutrients uptake and translocation in cucumber (*Cucumis sativus* L.) grown by the nutrient film technique. Ph. D thesis, London Univ., p: 299.
- El-Khateeb, M.A., Asmaa B. El-Attar and M. Rawda (2017). Application of Plant Bio stimulants to Improve the Biological Responses and essential oil production of marjoram (*Majorana hortensis*, Moench) plants. *Nour Middle East Journal of Agriculture Research* ISSN 2077-4605, **6(4)**: Pages: 928-941.
- El-Khateeb, M.A., A.B. El-Attar and R.M. Nour (2017). Application of plant bio stimulants to improve the biological responses and essential oil production of marjoram (*Majorana hortensis*, Moench) plants. *Middle East J. Agric. Res.*, **6(4)**: 928-941.
- El-Sherbeny, S.F., A.A. Hendawy, N.Y. Youssef and M.S. Hussein (2012). Response of turnip (*Brassica rapa*) plants to minerals or organic fertilizer treatments. *Journal of Applied Sciences Research*, **8(2)**: 628-634.
- Eqbal M.A.D. and A. Abdullah (2017). Medicinal and Functional Values of Thyme (*Thymus vulgaris* L.) Herb. *Journal of Applied Biology & Biotechnology*, **5(02)**: 017-022.
- Fachini-Queiroz, F.C., R. Kummer, C.F. Estevão-Silva, M.D.D.B. Carvalho, J.M. Da Cunha, R. Grespan, C.A. Bersani-Amado and R.K.N. Cuman (2012). Effects of Thymol and Carvacrol, Constituents of *Thymus vulgaris* L. Essential Oil, on the Inflammatory Response. *Evid. Based Complement. Altern. Med.*, 1-10.
- Fatma, S.A.M., A.M. Zeinab, M.A. Taha and U.A. El-Behairy (2008). Some chemical changes of *lavendula officinalis* plant treated with nicotine amid. *Biol. Chem. Environ. Sci.*, **3(2)**: 83-103.
- Fatma, S.A.M., A.M. Zeinab, Mohamed M. Shams El-Deen and U.A. El-Behairy (2014). Enhancement of physicochemical properties of strawberry produced under growth promoters and soilless conditions. the requirements for the degree of doctor of philosophy In Agricultural science (agricultural biochemistry) Department of agricultural biochemistry Faculty of agriculture Ain shams university.
- Fawzy, Z.F., M.M. El-magd, Y. Li, Z. Ouyang and A.M. Hoda (2012). Influence of foliar application by EM "effective microorganisms", amino acids and yeast on growth, yield and quality of two cultivars of onion plants under newly reclaimed soil. *J. Agric. Sci.*, **4**: 26-39.
- Fernanda, C.F.Q., K. Raquel, F.E.S. Camila, D.B.C. Maria, M. Joice, R.G. Cunha, A.B.A. Ciomar and K.N.C. Roberto (2012). Effects of Thymol and Carvacrol, Constituents of

- Thymus vulgaris* L. Essential Oil, on the Inflammatory Response. Volume 2012, Article ID 657026, 10 pages. doi: 10.1155/2012/657026.
- Gema, N. (2020). A Review on Applications and Uses of Thymus in the Food Industry, *Plants* 9, 961.
- Golkar, P., N. Mosavat and S.A.H. Jalali (2020). Essential oils, chemical constituents, antioxidant, antibacterial and in vitro cytotoxic activity of different Thymus species and Zataria multiflora collected from Iran. *S. Afr. J. Bot.*, **130**: 250–258.
- Gomez, M.L. and F.M. Lajolo (2008). Ascorbic acid metabolism in fruits: activity of enzymes involved in synthesis and degradation during ripening in mango and guava. *J. Sci. Food and Agric.*, **88**: 756-762.
- Guangzhe, Y., W. Qiuxing, H. Hao and X. Jixing (2020). Amino Acid Transporters in Plant Cells: A Brief Review. *Plants*, **9**: 967.
- Hanafy, M.S., A.A.M. Heikal, E.O. El-Ghawwas and Z.H. Ali (2017). The effect of cattle manure and biostimulants on growth and essential oil production of southernwood *Artemisia (Artemisia abrotanum L.)* plant. *Middle East J. Agric. Res.*, **6(4)**: 1218-1231.
- Hanafy, M.S., H.A. Ashour and F.M. Sedek (2018). Effect of some Bio-stimulants and Micronutrients on Growth, Yield and Essential Oil Production of *Majorana hortensis* plants. *International Journal of Environment*, ISSN 2077-4505 **7(1)**: Page: 37-52.
- Hashem, H.A.E. (2018). Response of Marjoram (*Majorana Hortensis L.*) Plant to foliar spraying by some antioxidants under Siwa Oasis conditions. *J. Agric. Vet. Sci.*, **11**: 51–58.
- Hassanain, M.A., E.M. Abdella and M.M. Rady (2006). Response of growth, flowering, oil yield and chemical composition of *Matricaria chamomilla L.* foliar applications. *J. Agric. and Env. Sci. Alex. Univ. Egypt*, **5(2)**: 33- 58.
- Heikal, A.E. (2005). Effect of organic and biofertilization on growth production and composition of (*Thymus vulgaris L.*) plants. M.Sc. Thesis, Fac. Agric. Cairo Univ.
- Hemdan, S.H. (2008). Effect of some organic and biofertilization treatments on anise plants. M.Sc. Thesis, Fac. of Agric., Minia University.
- Hendawy, S.F., Azza A Ezz El-Din, Eman E. Aziz and E.A. Omer (2010). Productivity and oil quality of *Thymus vulgaris L.* under organic fertilization conditions. *Ozean Journal of Applied Sciences*, **3(2)**: 203-216.
- Hendawy, S.F., M.S. Hussein, A.E. El-Gohary and M.E. Ibrahim (2015). Effect of Foliar Organic Fertilization on the Growth, Yield and Oil Content of *Mentha piperita var. citrata*. *Asian Journal of Agricultural Research*, **9(5)**: 237-248.
- Kamal, H.M., M.A. Elisa and A.A. Mohammed (2017). 'Effect of some mineral compounds on yield and fruit quality of pomegranate', *Bioscience Research*, **14(4)**: pp. 1197-1203.
- Kenawy, A.G. (2010). Effect of some biofertilization treatments on the growth, yield and chemical composition of *Hibiscus sabdariffa L.* plants. M.Sc. Thesis, Fac. of Agric., Mania Univ.
- Khattab, M.M., A.E. Shaban, A.H. El-Shrief and A.S. Mohamed (2012). Effect of humic Acid and Amino Acids on Pomegranate Trees under Deficit Irrigation. I: Growth, Flowering and Fruiting. *Journal of Horticultural Science and Ornamental Plants*, **4(3)**: pp. 253-259.
- Kocira S. (2019). Effect of amino acid bio stimulant on the yield and nutraceutical potential of soybean. *Chilean journal of agricultural research*, **79(1)**: january-march 2019.
- Maher, A.A., S.M. Al. Maqtari and Alghalibi E.H. Alhamzy (2011). Chemical composition and antimicrobial activity of essential oil of *Thymus vulgaris* from Yemen. *Turk. J. Biochem.*, **36**: 342-349.
- Mahmoud, H.I., N.A. Azzaz, Y.A.M. Khalifa, M.A. Mahmoud and G. Fakhry (2016). Effect of foliar application with active yeast extract and benzyladenine on some vegetative growth criteria and chemical composition of lupine (*Lupinus termis L.*) plants. *Minia J. Agric. Res. Develop.*, **36**: 193-214.
- Matter, F.M.A. and S.A.A. El Sayed (2015). Influence of mixed NPK fertilizers with foliar spray of active dry yeast on caraway under newly reclaimed soil conditions. **5(2)**: 423-430.
- Mohamed, A.K.A., H.A. Abdel-Galil and H.G. Naglaa (2020). Effect of some nutrients and amino acids spraying on yield and fruit quality of Manfalouty pomegranate. *S/U-International Journal of Agricultural Science*, **2(2)**: pp.: 18-29.
- Mohamed, M.H., A.B. Elham, Sh.S. Mervat and H.K. Howida (2020). Effect of garlic extract, ascorbic acid and nicotinamide on growth, some biochemical aspects, yield and its components of three faba bean (*Vicia faba L.*) cultivars under sandy soil conditions.
- Mohamed, S.M., E.M. Abou El-Ghait, N.S.A. El Shayeb, Y.A. Ghatas and A.A. Shahin (2015). Effect of some fertilizers on improving growth and oil productivity of basil (*Ocimum basilicum L.*) cv. Genovese plant. *Egypt. J. Appl. Sci.*, **30(6)**: 384-399.
- Montagne, V., S. Charpentier, P. Cannavo, H. Capiiaux, C. Grosbellet and T. Lebeau (2015). Structure and activity of spontaneous fungal communities in organic substrates used for soilless crops. *Sci. Hort.*, **192**: 148-157.
- Murphy, R.J. and J.P. Riley (1962). A modified single solution method for determination of phosphate in natural waters. *Anal. Chim. Acta.*, **27**: 31-36.
- Naglaa, M.M. (2020). Effect of foliar application of some vitamins and irrigation intervals on vegetative growth, flowering and some biochemical constituents of *Helianthus annuus L.* plants. *Scientific Journal of Agricultural Sciences*, **2(2)**: 1-8.
- Nahed, G.A.A., A.M.M. Azza and M.M. Farahat (2010). Response of vegetative growth and chemical constituents

- of *Thuja orientalis* L. plant to foliar application of different amino acids at Nubaria. *J. Am. Sci.*, **6(3)**: 295- 301.
- Olga, K., Bc. Katarzyna, L.P. Jaroslaw, P. Anna, R. Katarzyn and W. Zenon (2020). Morphological and Chemical Traits as Quality Determinants of Common Thyme (*Thymus vulgaris* L.), on the Example of 'Standard Winter' Cultivar. *Agronomy*, **10**: 909.
- Prabodh, S.B.I., R. Burray, S. Mcfeeter and N.S. William (2016). Essential oil characterization of thymus vulgaris from various geographical locations. *Foods*, **5**: 70.
- Prasanth, R.V., V.K. Ravi, P.V. Varsha and S. Satyam (2014). Review on *Thymus vulgaris* Traditional Uses and Pharmacological Properties. *Med. Aromat Plants*, **3**: 164.
- Putra, P.A. and H. Yuliando (2015). Soilless culture system to support water use efficiency and product quality: A review. *Agr. Agr. Sci. Procedia.*, **3**: 283-288.
- RCT (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Radwan, S.S. (1978). Coupling of two dimension thin layer chromatography with gas chromatography for the quantitative analysis of lipids classes and their constituent fatty acids. *J. Chromatogram Sci.*, **16**: 538-542.
- Refaat, A.M. and L.K. Balbaa (2001). Yield and quality of lemongrass plants (*Cymbopogon flexuosus* Stapf) in relation to foliar application of some vitamins and microelements. *Egypt. J. Hort.*, **28(1)**: 41-57.
- Roby, M.H.H., M.A. Sarhan, K.A.H. Selim and K.I. Khalel (2013). Evaluation of antioxidant activity, total phenols and phenolic compounds in thyme (*Thymus vulgaris* L.), sage (*Salvia officinalis* L.) and marjoram (*Origanum majorana* L.) extracts. *Ind. Crop. Prod.*, **48**: 43-48.
- Sarikurkcuc, C., G. Zengin, M. Oskay, S. Uysal, R. Ceylan and A. Aktumsek (2015). Composition, antioxidant, antimicrobial and enzyme inhibition activities of two *Origanum vulgare* subspecies (subsp. *vulgare* and subsp. *hirtum*) essential oils. *Ind. Crops Prod.*, **70**: 178-184.
- Shadia, K.A., Kh.A. Hammam and A.A. Alia (2014). Effect of bio- fertilization and some plant extracts on the growth, yield and chemical constituents of basil plant. *J. Plant Production, Mansoura Univ.*, **5(2)**: 193- 210.
- Shafeek, M.R., Y.I. Helmy, M.M. Neama, A.F.Sh. Magda and M.O. Nadia (2013). Effect of foliar application of some antioxidants on growth, yield and chemical composition of Lettuce plants (*Lactuca Sativa* L.) under plastic house condition. *Middle East Journal of Applied Sciences*, **3(2)**: 70-75.
- Shahad, A., H. Zainab, M. Wafa, Amal Abdullah and A.K. Shah (2020). Chemical composition, in vitro antibacterial and antioxidant potential of Omani Thyme essential oil along with in silico studies of its major constituent. *Journal of King Saud University – Science*, **32**: 1021-1028.
- Shahidi, F. and M. Naczk (1995). Methods of Analysis and Quantification of Phenolic Compounds. Food phenolic: sources, chemistry, effects and applications (pp. 287-293). Techno. Pub. Company, Inc: Lancaster.
- Sharangi, A.B. and S. Guha (2013). Wonders of leafy spices: Medicinal properties ensuring Human Health. *Science International*, 312-317.
- Shinwari, Z. (2005). Medicinal plants research in the 21st. century. International symposium Medicinal Plants: Linkages beyond national boundaries., (Eds) Z. Shinwari, T. Watanabe and M.Ali, pp.12-16.
- Wanas, A.L. (2006). Trails for improving growth and productivity of tomato plants grown in winter. *Annals. Agric. Sci. Moshtohor*, **44(3)**: 466-471.
- Wassel, A.H.M., A.A. Gobara, H.I.M. Ibrahiem and M.M. Shaaban (2015). Response of Wonderful Pomegranate trees to foliar application of Amino Acids, Vitamins B and Silicon. *World Rural Observations*, **7(3)**: pp. 91-95.
- Watanabe, F.S. and S.R. Olsen (1965). Test of an ascorbic acid method for determining phosphorous in water and Na HCO₃ extracts from Soil. *Soil sci. Soc. Am. Proc.*, **29**: 677-678.
- Wettstein, D.V. (1957). Chlorophyll- Lethale und der submikroskopische Formwechsel der Plastiden. *Exp. Cell Res.*, **12**: 427-506.