



## EFFECT OF INTEGRATED NUTRIENT AND WEED MANAGEMENT PRACTICES ON WEEDS, GROWTH AND YIELD OF SUNFLOWER

Kalaiyarasan, C. and V. Vaiyapuri

Department of Agronomy, Faculty of Agriculture,  
Annamalai University, Annamalai Nagar – 608 002, Tamil Nadu, India.  
Corresponding Author: kalai77.agri@gmail.com

### Abstract

A comprehensive study was made in irrigated sunflower to optimise the integrated nutrient and weed management practices for augmenting sunflower productivity, at Annamalai University Experimental Farm, Annamalai Nagar, Tamilnadu, India. The experiment was laid out in split plot design with three replications. The details of the treatment in mainplots are M<sub>1</sub>-Control, M<sub>2</sub>-RDF (40:20:20 kg NPK ha<sup>-1</sup>) + FYM @ 12.5 t ha<sup>-1</sup>, M<sub>3</sub>-RDF+ Vermicompost @ 5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> + foliar spray of 1% KH<sub>2</sub>PO<sub>4</sub> (twice at 25 and 55 DAS), M<sub>4</sub>-RDF+ FYM @ 12.5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> + foliar spray of 1% KH<sub>2</sub>PO<sub>4</sub> (twice at 25 and 55 DAS) M<sub>5</sub>-RDF+ Vermicompost @ 5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup>, M<sub>6</sub>-RDF+ FYM @ 12.5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> + foliar spray of 1% KH<sub>2</sub>PO<sub>4</sub> (twice at 25 and 55 DAS) and the subplots are S<sub>1</sub>- Unweeded control, S<sub>2</sub>- Pre emg. Oxyflourfen @ 0.1 kg ha<sup>-1</sup> + HW at 30 DAS, S<sub>3</sub>- Pre sowing fluchloralin @ 1 kg ai ha<sup>-1</sup> + HW at 30 DAS, S<sub>4</sub>- Pre emg. Pendimethalin @ 1 kg ai ha<sup>-1</sup> + HW at 30 DAS and S<sub>5</sub>- HW twice at 15 and 30 DAS. The results of the study evidently proved that application of recommended NPK+ vermicompost + Azospirillum+ ZnSO<sub>4</sub> + foliar spray of KH<sub>2</sub>PO<sub>4</sub> along with fluchloralin + HW at 30 DAS (M<sub>3</sub>S<sub>5</sub>) as an agronomically efficient, eco-friendly and economically viable technology for improving sunflower yield and quality. This treatment (M<sub>3</sub>S<sub>5</sub>) combination registered lowest values for weed density, nutrient removal by weeds, weed biomass and maximum weed control index and maximum values for growth and yield attributes and yield of sunflower in both the crops.

**Key words** : Integrated nutrient, Sunflower etc.

### Introduction

The cultivated sunflower (*Helianthus annus* L.) is an annual oilseed plant of *compositae* family. Sunflower has many advantages over other oilseeds crops. The crop is endowed with short growth period, photosensitiveness and presence of high degree of polyunsaturated fatty acid (PUFA) content. The sunflower oil has a pleasant flavour and excellent keeping quality when refined. Cholesterol lowering factor constitutes around 85-90% of the total fatty acid (Silver *et al.*, 1984). Fertilizer application as the major input through which the productivity can be increased by exploiting varietal potential. Chemical fertilizers have had a substantial impact on yield increments in the recent past and are today an indispensable part of modern agricultural practices (Reddy and Raja Reddy, 2002). Integration of organic manures and biofertilizers with chemical fertilizers is more emphasised not only to boost the production of sunflower from limited land resource but also for its sustainability. There is need to promote use of organics in addition to inorganic fertilizers for sustained maintenance of soil fertility (Devidayal and Agarwal, 1999). Sunflower which grows slowly during its initial stage provides congenial

environment for weed growth in abundance. The weeds cause drastic reduction in seed yield of sunflower upto 83% (Legha *et al.*, 1992). The critical period of weed competition is upto 30 DAS in sunflower (Muthusankaranarayan *et al.*, 1995). The most promising single approach to weed control in land reported is to combine manual, cultural and mechanical methods with herbicides (Yaduraju and Mishra, 2003).

### Material and Methods

The field experiments were conducted to study the effect of integrated nutrient and weed management on sunflower at Experimental Farm, Department of Agronomy, Annamalai University, Annamalai Nagar (TN). The soil of experimental field was clayey loam with low in available nitrogen (212.4 kg ha<sup>-1</sup>), medium in available phosphorus (28.3 kg ha<sup>-1</sup>) and high in available potassium (348.1 kg ha<sup>-1</sup>). The pH and E.C. were 7.5 and 0.45 dsm<sup>-1</sup> respectively. The experiment was laid out in a split plot design with three replication. The details of the treatment in mainplots are M<sub>1</sub>-Control, M<sub>2</sub>-RDF (40:20:20 kg ha<sup>-1</sup>) + FYM @ 12.5 t ha<sup>-1</sup>, M<sub>3</sub>-RDF + Vermicompost @ 5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> + foliar spray of 1% KH<sub>2</sub>PO<sub>4</sub> (twice at 25 and 55

DAS), M<sub>4</sub>-RDF+ FYM @ 12.5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> + foliar spray of 1% KH<sub>2</sub>PO<sub>4</sub> (twice at 25 and 55 DAS) M<sub>5</sub>- RDF+ Vermicompost @ 5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup>, M<sub>6</sub>- RDF+ FYM @ 12.5 t ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> + seed treatment with Azospirillum (600 g ha<sup>-1</sup>) + ZnSO<sub>4</sub> @ 25 kg ha<sup>-1</sup> and the subplots are S<sub>1</sub>- Unweeded control, S<sub>2</sub>- Pre emg. Oxyflourfen @ 0.1 kg ha<sup>-1</sup> + HW at 30 DAS, S<sub>3</sub>- Pre sowing fluchloralin @ 1 kg ai ha<sup>-1</sup> + HW at 30 DAS, S<sub>4</sub>- Pre emg. Pendimethalin @ 1 kg ai ha<sup>-1</sup> + HW at 30 DAS, S<sub>5</sub>- HW twice at 15 and 30 DAS. Recommended dose of 40:20:20 kg of NPK ha<sup>-1</sup> was applied. N was applied in the form of urea while phosphorus and potassium were applied in the form of SSP and MOP respectively. Entire dose of P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O and half of N was applied as basal and remaining "N" at 30 DAS. Weed management practices were carried out as per the treatment schedule. The pre emergence herbicides (Pendimethalin, oxyflourfen and metalachlor) at required quantities were taken and sprayed at 3 DAS using the hand operated knapsack sprayer fitted with a flood a jet nozzle. A spray volume of 500 litres of water was used per hectare.

## Results and Discussion

### Weeds (Table 1 and 2)

The nutrient management treatments significantly influenced the weed characters in sunflower. Among the nutrient management practices tried, the treatment M<sub>3</sub> (RDF+ vermicompost + azospirillum + ZnSO<sub>4</sub> + KH<sub>2</sub>PO<sub>4</sub>) recorded lower weed population (378.20 and 448.60 m<sup>-2</sup>) and (390.00 and 462.00m<sup>-2</sup>), lesser weed biomass (97.23 and 107.26 kg ha<sup>-1</sup>) and (104.34 and 102.47 kg ha<sup>-1</sup>), higher weed control index (77.01 and 80.45 %) and (76.63 and 81.73 %) at 15 and 30 DAS in first and second crop respectively. This treatment also record lesser nitrogen removal by weeds (16.10 and 17.20 kg ha<sup>-1</sup>), phosphorus removal by weeds (4.03 and 4.20 kg ha<sup>-1</sup>), potassium removal by weeds (13.44 and 12.56 kg ha<sup>-1</sup>) at 30 DAS in first and second crop respectively. The reason for low weed population under these treatments might be due to better uptake of nutrients by the crop from the initial stage and did not provide enough time for the weeds to utilise the nutrients and other factors. Similar result was reported by Patel *et al.* (1995). This was followed by M<sub>4</sub> (RDF+FYM+Azospirillum+ZnSO<sub>4</sub>+KH<sub>2</sub>PO<sub>4</sub>). Highest values for weed density, weed biomass and nutrient removal were recorded in M<sub>1</sub> (No NPK/ Organics).

Profound influence on weed count was noticed due to weed management treatments. Among the different weed management practices tried, S<sub>3</sub> (fluchloralin +

HW at 30 DAS) registered the lowest weed count (263.83 and 279.16 m<sup>-2</sup>) and (338.5 and 350.00m<sup>-2</sup>), lowest weed biomass (89.12 and 95.01 kg ha<sup>-1</sup>) and (79.21 and 85.2kg ha<sup>-1</sup>), highest weed control index (78.93 and 78.72%) and ( 85.56 and 81.79 %)at 15 and 30 DAS in first and second crop respectively. S<sub>5</sub> (HW twice at 15 and 30 DAS) recorded a lesser nutrient removal nitrogen removal by weeds (14.54 and 15.54 kg ha<sup>-1</sup>), phosphorus removal by weeds (3.70 and 3.70 kg ha<sup>-1</sup>) and potassium removal by weeds (13.52 and 12.62 kg ha<sup>-1</sup>) at 30 DAS in first and second crop respectively. It may be due to the efficiency of the sowing herbicide in supporting germination of weed seeds. This findings is in conformity with the studies of Rodrique *et al.* (1982). The unweeded control (S<sub>1</sub>) treatment recorded higher weed density, weed biomass, poor weed and maximum NPK removal the crops at all the stages. This is due to poor weed management.

Significant interactions were noticed between the nutrient and weed management practices in both the crops. The Interaction between nutrient management (M<sub>3</sub>) with the weed management treatment (S<sub>5</sub>) proved efficiency by registering lowest weed density, biomass, nutrient removal by weeds and maximum weed control index . This might be due to the herbicidal effect of fluchloralin might be due to the inhibition of cell division through tubulin inactivation mechanism which might have curtailed the density and growth of weeds Krishne Gowda *et al.* (1985).

### Crop Growth Attributes (Table 3)

Among the nutrient management practices tried, the treatment M<sub>3</sub> (RDF + vermicompost + azospirillum + ZnSO<sub>4</sub> + KH<sub>2</sub>PO<sub>4</sub>) recorded maximum plant height (145.34 cm) at harvest stage, leaf area index (6.46) at flowering stage and dry matter production (4449.13 kg ha<sup>-1</sup>) at harvest stage, root length (27.92cm), root volume (17.20 cm<sup>-3</sup> plant<sup>-1</sup>). Lowest plant height, leaf area index and dry matter production recorded under M<sub>1</sub> (control) in all stages of crop growth.

Among the weed management treatments, S<sub>5</sub> (HW twice at 30 DAS) recorded maximum plant height (141.37 cm) at harvest stage, leaf area index (6.22) at flowering stage and dry matter production (4006.97 kg ha<sup>-1</sup>) at harvest stage, root length (27.02cm), root volume (16.7cm<sup>-3</sup> plant<sup>-1</sup>). This was followed by the treatment S<sub>3</sub> (fluchloralin + HW at 30 DAS). The plant height, leaf area index and dry matter production recorded under S<sub>1</sub> (unweeded control) in all stages of crop growth.

The Interaction effect between the nutrient and weed management on plant growth attributes is significant. Treatment M<sub>3</sub> (RDF + vermicompost + azospirillum + ZnSO<sub>4</sub> + KH<sub>2</sub>PO<sub>4</sub>) with S<sub>5</sub> (HW twice

at 30 DAS) maximum plant height, leaf area index (7.35) at flowering stage and dry matter production (4521.13 kg ha<sup>-1</sup>) at harvest stage, root length (31.2cm), root volume (18.6 cm<sup>3</sup> plant<sup>-1</sup>). Lowest plant height, leaf area index and dry matter production recorded under M<sub>1</sub>S<sub>1</sub> (control) in all stages of crop growth.

This might be due to the effective interaction between the nutrient and weed management treatments, which could have increased the availability of better nutrition from vermicompost and other components along with the efficient control of weeds by the respective treatments. Similar trend of results was reported by Patel *et al.* (1994).

#### Yield Attributes (Table 4 and 5)

Among the nutrient management practices tried M<sub>3</sub> (RDF+ vermicompost+ azospirillum+ ZnSO<sub>4</sub>+ KH<sub>2</sub>PO<sub>4</sub>) recorded maximum values for head diameter (18.5cm), total number of seeds head<sup>-1</sup> (866.2head<sup>-1</sup>), number of filled seeds head<sup>-1</sup>(513.7), seed filling percentage (94.8), test weight(7.73g), seed yield (1671kg ha<sup>-1</sup>) and stalk yield (5752kg ha<sup>-1</sup>) over other treatments. This was followed by M<sub>4</sub>(RDF + FYM + azospirillum + ZnSO<sub>4</sub> + KH<sub>2</sub>PO<sub>4</sub>). M<sub>1</sub> (control) recorded lower value for head diameter (14.03cm), total number of seeds head<sup>-1</sup> (827.18.head<sup>-1</sup>), number of filled seeds head<sup>-1</sup> (466.22), seed filling percentage, test weight(6.10g), seed yield (503kg ha<sup>-1</sup>) and stalk yield. Among the weed

management treatments S<sub>5</sub> (HW twice at 30 DAS) registered higher head diameter (18.7cm), total number of seeds head<sup>-1</sup> (837.4 head<sup>-1</sup>), number of filled seeds head<sup>-1</sup> (786.4), seed filling percentage (93.5), test weight(7.60g), seed yield (1201kg ha<sup>-1</sup>) and stalk yield (5622kg ha<sup>-1</sup>) over other treatments. This was followed by S<sub>3</sub> (fluchloralin + HW at 30 DAS). unweeded control (S<sub>1</sub>) recorded lowest head diameter, total number of seeds head<sup>-1</sup>, number of filled seeds head<sup>-1</sup>, seed filling percentage, test weight, seed yield and stalk yield.

The Interaction effect between the nutrient and weed management was significant. Treatment M<sub>3</sub> (RDF + vermicompost + azospirillum+ZnSO<sub>4</sub>+KH<sub>2</sub>PO<sub>4</sub>) with S<sub>5</sub> (HW twice at 30 DAS) registered higher head diameter (20.31cm),total number of seeds head<sup>-1</sup> (946.21 head<sup>-1</sup>), number of filled seeds head<sup>-1</sup>(929.25), seed filling percentage, test weight(8.13g), seed yield (1901kg ha<sup>-1</sup>) and stalk yield (6225kg ha<sup>-1</sup>) over other treatments. This was followed by M<sub>3</sub>S<sub>3</sub> and lowest yield was recorded by M<sub>1</sub>S<sub>1</sub> head diameter, total number of seeds head<sup>-1</sup>, number of filled seeds head<sup>-1</sup>, seed filling percentage, test weight, seed yield and stalk yield.

These findings are in conformity with the findings of Babusaravanan (1992) in groundnut. These results indicated that integrated nutrient management under comparatively weed free environment can influence the sunflower yield components and seed yield significantly.

**Table 1:** Effect of integrated nutrient and weed management practices on weed characters of sunflower

Treatments	Mean Weed Population (M <sup>2</sup> )				Weed Biomass (Kg ha <sup>-1</sup> )			
	I Crop		II Crop		I Crop		II Crop	
	15 DAS	30 DAS	15 DAS	30 DAS	15 DAS	30 DAS	15 DAS	30 DAS
M <sub>1</sub>	417.00 (20.29)	529.60 (22.84)	448.40 (21.03)	543.40 (23.14)	387.7	435.7	384.8	436.5
M <sub>2</sub>	403.60 (19.95)	516.00 (22.54)	420.80 (20.38)	532.20 (22.89)	345.9	386.6	343.1	388.25
M <sub>3</sub>	378.20 (19.25)	448.60 (20.97)	390.00 (19.56)	462.00 (21.30)	97.23	107.3	104.3	102.5
M <sub>4</sub>	386.00 (19.47)	456.60 (21.18)	401.00 (19.86)	469.00 (21.47)	134.5	115.9	116.6	114.6
M <sub>5</sub>	394.00 (19.69)	486.20 (21.84)	411.00 (20.12)	495.20 (22.04)	303.2	356.6	317.9	360.6
M <sub>6</sub>	397.40 (19.79)	503.40 (22.23)	414.80 (20.23)	517.80 (22.56)	328.3	374.4	327.8	367.9
SEd	0.038	0.41	0.55	0.05	3.90	1.95	4.88	3.96
CD (P=0.05)	0.08	0.093	0.12	0.12	7.85	3.92	7.70	7.96
Sub Plot								
S <sub>1</sub>	505.83 (22.49)	695.16 (26.36)	524.16 (22.90)	710.66 (26.55)	422.9	548.73	446.7	560.9
S <sub>2</sub>	393.00 (19.81)	573.66 (23.94)	422.16 (20.50)	588.66 (24.25)	291.9	416.87	301.3	412.9
S <sub>3</sub>	263.81 (16.24)	373.33 (19.50)	279.16 (16.71)	386.00 (19.63)	89.1	108.9	95.0	96.4
S <sub>4</sub>	310.83 (17.64)	469.66 (21.67)	326.33 (18.07)	481.00 (21.93)	180.6	326.6	165.2	319.8
S <sub>5</sub>	506.66 (22.51)	338.50 (18.39)	519.83 (22.80)	350.00 (18.70)	346.0	79.2	307.2	85.3
S.Ed	0.05	0.05	0.073	0.005	2.76	1.38	3.45	3.26
CD (P=0.05)	0.101	0.11	0.14	0.011	6.16	3.08	9.82	

• : Figures in parenthesis are arc sin transformed values.

**Table 2:** Effect of integrated nutrient and weed management practices on weed control index (WCI) and Nutrient removal by weeds on Sunflower

Treatments	WCI (%)				Nutrient removal by weeds (kg ha <sup>-1</sup> ) at 30 DAS					
	I Crop		II Crop		I Crop			II Crop		
Main plot	15	30 DAS	15	30 DAS	N	P	K	N	P	K
M <sub>1</sub>	8.33	20.60	13.85	22.18	22.7	5.51	19.94	23.9	5.73	18.5
M <sub>2</sub>	18.22	29.55	23.19	30.77	18.1	4.68	16.93	19.5	4.64	15.64
M <sub>3</sub>	77.01	80.45	76.63	81.73	16.1	4.03	13.44	17.2	4.20	12.6
M <sub>4</sub>	68.19	78.87	73.90	79.56	16.7	4.19	15.78	17.8	4.35	14.8
M <sub>5</sub>	28.31	35.01	28.83	35.76	17.0	4.29	15.86	18.2	4.42	14.9
M <sub>6</sub>	22.38	31.76	26.61	34.40	17.4	4.43	16.06	18.8	4.50	15.02
S.Ed					0.078	0.09	0.0039	0.043	0.011	0.017
CD (P=0.05)					0.157	0.019	0.007	0.086	0.023	0.0035
Sub Plot										
S <sub>1</sub>	-	-	-	-	28.7	6.72	25.09	29.9	6.86	23.7
S <sub>2</sub>	30.97	24.03	32.55	26.37	17.1	4.56	15.45	18.7	4.66	14.2
S <sub>3</sub>	78.93	30.16	78.72	82.81	14.8	3.76	13.72	15.9	3.92	12.8
S <sub>4</sub>	57.29	41.047	63.01	42.97	15.0	3.85	13.90	16.1	4.0	12.9
S <sub>5</sub>	18.19	85.56	31.22	84.79	14.5	3.70	13.52	15.5	3.76	12.6
S.Ed					0.055	0.006	0.0027	0.030	0.008	0.0012
CD (P=0.05)					0.123	0.015	0.006	0.067	0.018	0.0027

**Table 3:** Effect of integrated nutrient and weed management practices on growth attributes of sunflower

Treatments	Plant height (cm) (At harvest)		LAI (At flowering)		DMP (Kg ha <sup>-1</sup> ) (At harvest)		Root length (cm) (At 60 DAS)		Root volume (Cm <sup>3</sup> / plant) (At 60 DAS)	
	I	II	I	II	I	II	I	II	I	II
Main plot										
M <sub>1</sub>	103.0	79.9	4.15	4.06	3297	2954	20.5	18.2	13.7	12.9
M <sub>2</sub>	125.8	105.0	5.41	5.28	3958	3637	25.1	22.8	15.9	15.4
M <sub>3</sub>	145.3	124.9	6.46	6.31	4449	4103	27.9	26.2	17.2	16.9
M <sub>4</sub>	138.6	118.6	6.10	6.03	4291	3953	26.4	24.5	16.6	16.4
M <sub>5</sub>	135.6	116.2	5.95	5.88	4230	3898	26.0	24.2	16.4	16.2
M <sub>6</sub>	131.7	112.1	5.75	5.65	4099	3756	25.6	23.9	16.2	15.9
S.Ed	0.409	0.37	0.002	0.003	14.9	16.3	0.19	0.15	0.043	0.048
CD (P=0.05)	0.91	0.84	0.051	0.01	29.8	32.7	0.39	0.32	0.088	0.098
Sub Plot										
S <sub>1</sub>	111.8	88.9	4.69	4.64	3509	3250	22.3	20.8	14.7	14.1
S <sub>2</sub>	121.9	101.7	5.27	5.15	3848	3481	24.2	21.9	15.4	15.1
S <sub>3</sub>	139.1	118.9	6.08	5.98	4219	3951	26.5	24.8	16.7	16.3
S <sub>4</sub>	135.9	116.4	5.93	5.81	4220	3879	26.2	24.4	16.5	16.1
S <sub>5</sub>	141.4	121.3	6.22	6.09	4402	4006	27.0	25.4	16.9	16.5
S.Ed	0.213	0.07	0.018	0.003	12.7	14.1	0.20	0.09	0.036	0.039
CD (P=0.05)	0.42	0.15	0.037	0.001	25.5	28.3	0.40	0.19	0.073	0.079

**Table 4:** Effect of integrated nutrient and weed management practices on yield attributes of sunflower

Treatments	50% flowering		Head diameter (cm)		Total no. of seeds head <sup>-1</sup>		Number of filled Seeds head <sup>-1</sup>		Seed filling (%)		Test Wt. (g)	
	I	II	I	II	I	II	I	II	I	II	I	II
<b>Main Plot</b>												
M <sub>1</sub>	56.0	58.5	14.0	13.8	578.7	479.8	466.2	365.6	79.5	78.8	6.10	6.07
M <sub>2</sub>	51.8	52.8	16.4	16.2	753.7	643.5	683.5	574.7	90.4	89.1	7.29	7.28
M <sub>3</sub>	50.2	50.9	18.5	18.2	866.2	774.0	513.7	721.2	94.8	93.7	7.73	7.70
M <sub>4</sub>	50.8	51.5	18.1	17.8	826.1	734.3	770.9	676.5	93.0	91.8	7.58	7.56
M <sub>5</sub>	51.0	51.7	17.8	17.4	814.0	718.7	753.2	651.8	92.6	91.3	7.51	7.48
M <sub>6</sub>	51.4	52.3	17.2	17.0	785.6	678.9	723.7	614.4	92.0	90.1	7.41	7.39
S.Ed	0.25	0.029	0.005	0.0057	3.82	3.44	2.29	1.48	0.058	0.054	0.020	0.019
CD (P=0.05)	0.51	0.06	0.0112	0.0166	8.53	6.92	4.61	2.98	0.126	0.109	0.041	0.04
<b>Sub Plot</b>												
S <sub>1</sub>	53.9	55.9	14.7	14.6	648.3	546.4	544.3	453.5	83.0	82.1	6.60	6.58
S <sub>2</sub>	52.3	53.8	15.8	15.6	727.3	622.7	653.7	544.9	89.7	88.0	7.13	7.10
S <sub>3</sub>	50.9	51.9	18.41	17.9	827.2	731.4	772.2	676.8	93.2	91.9	7.54	7.52
S <sub>4</sub>	51.3	51.9	18.44	17.6	814.1	712.2	752.9	653.5	92.5	91.3	7.47	7.46
S <sub>5</sub>	50.7	51.4	18.7	18.1	837.4	744.9	786.4	691.4	93.5	92.2	7.60	7.58
S.Ed	0.19	0.024	0.004	0.0056	3.53	2.94	0.129	1.21	0.056	0.046	0.016	0.014
CD (P=0.05)	0.39	0.05	0.009	0.0114	7.1	6.55	2.61	2.43	0.118	0.093	0.033	0.03

**Table 5:** Effect of integrated nutrient and weed management practices on yield and quality of sunflower

Treatments	Seed yield (Kg ha <sup>-1</sup> )		Stalk yield (Kg ha <sup>-1</sup> )		Oil Content (%)		Protein Content %	
	I	II	I	II	I	II	I	II
<b>Main Plot</b>								
M <sub>1</sub>	503	489	4279	4121	37.30	37.31	18.26	17.14
M <sub>2</sub>	826	817	5160	5054	38.48	38.34	18.63	17.61
M <sub>3</sub>	1671	1591	5752	5536	39.18	39.03	18.82	18.04
M <sub>4</sub>	1263	1212	5550	5368	38.83	38.81	18.75	17.90
M <sub>5</sub>	182	1085	5471	5311	38.73	38.68	18.73	17.85
M <sub>6</sub>	988	979	5326	5187	38.63	38.50	18.69	17.76
S.Ed	22.96	16.37	17.46	83.94	0.0069	0.0029	0.0079	0.003
CD (P=0.05)	46.24	32.91	34.97	19.93	0.014	0.006	0.016	0.006
<b>Sub Plot</b>								
S <sub>1</sub>	833	801	4644	4544	37.79	37.74	18.45	17.27
S <sub>2</sub>	1009	967	4987	4903	38.29	38.33	18.54	17.55
S <sub>3</sub>	1169	1128	5546	5352	38.83	38.76	18.76	17.92
S <sub>4</sub>	1116	1088	5483	5278	38.76	38.66	18.72	17.87
S <sub>5</sub>	1201	1161	5622	5404	38.95	38.74	18.78	17.96
S.Ed	15.57	15.35	15.94	17.40	0.0059	0.0019	0.071	0.002
CD (P=0.05)	31.31	30.86	31.92	34.99	0.012	0.004	0.014	0.004

**References**

- Babusaravanan, K. (1992). A study on methods of pendimethalin application to groundnut at different nitrogen regimes. M.Sc.(Ag.) Thesis, AC & RI, TNAU, Mdurai.
- Devi, D. and Agarwal, S.K. (1998). Response of sunflower (*Helianthus annuus* L.) to organic manures and fertilizers. Indian J. Agron., 43(3): 469-473.
- Krishne Gowda, K.T.; Muniyappa, T.V. and Venkatramu, M.N. (1985). Weed control in sunflower through mechanical, chemical and cultural method. Indian J. Weed Sci., 17(4): 49-51.
- Legha, P.K.; Malik, R.K. and Faroda, A.S. (1992). Weed management in Kharif sunflower. Crop Res., 5(2): 376-378.
- Muthusankaranarayan, A.; Veerabadran, V. and Balasubramanian, R. (1995). Critical period of crop-weed competition in rainfed sunflower. Abstract of paper VI Biennial Conference on Indian Society of Weed Science. pp. 67.
- Patel, Z.G.; Parmer, N.D. and Raj, V.C. (1994). Effect of weed control methods on yield and yield attributes of sunflower. Indian J. Agron., 39(2): 330-331.
- Patel, D.M.; Patel, S.K. and Karelia (1995). Integrated nutrient management in pearl millet (*Pennisetum glaucum*) – wheat (*Triticum aestivum*) cropping system. Indian J. Agron., 40(2): 266-268.
- Reddy, P.V.R.M and Raja Reddy, C.K. (2002). Yield and yield attributes of sunflower as influenced by integrated nutrient management. The Andhra Agric. J., 49(1&2): 41- 44.
- Silver, J.G.; Rochester, C.P.; Bishop, D.G. and Harris, H.C. (1987). Unsaturated fatty acid synthesis during the development of isolated sunflower seeds. J. Exp. Bot., 35(1580): 1507-1514.
- Yaduraja, N.T. and Mishra, J.S. (2003). Weed management in Oilseed crops. In: proceedings of national seminar on stress management in oilseeds for attaining self-reference in vegetable oils, Directorate of Oilseeds Research, Jan. 28-30, Hyderabad, India, 49-71.