



## STUDIES ON THE EFFECT OF ORGANIC MANURES, BIOSTIMULANTS AND MICRONUTRIENTS ON CERTAIN GROWTH AND FLOWERING PARAMETERS OF TUBEROSE (*Polianthes tuberosa* L.) CV. PRAJWAL

R. Sureshkumar\*\*, G. Sahana Priya\*, M. Rajkumar\*\* and R. Sendhilnathan\*\*

\*PG Scholar, Department of Horticulture

\*\*Assistant Professor, Department of Horticulture

Annamalai University, Tamil Nadu-608 002.

Corresponding author E-mail ID: hortsuresh99@gmail.com

### Abstract

The present investigation entitled “studies on the effect of organic manures, biostimulants and micronutrients on certain growth and flowering parameters of tuberose (*Polianthes tuberosa* L.) Cv. Prajwal” was carried out in the Department of Horticulture, Faculty of Agriculture, Annamalai University, Annamalainagar during 2017-2018. The different treatments included the combinations of vermicompost, poultry manure, neem cake, humic acid, seaweed extract, zinc and boron in different ratios. The data was analyzed statistically which showed significant effect of organic manures, biostimulants and micronutrients over control values. The maximum values of plant height, number of leaves plant<sup>-1</sup> and number of sideshoots plant<sup>-1</sup>, spike length, flower diameter, rachis length and number of floret spike<sup>-1</sup> were observed in 50% RDF @ 100: 100: 100 kg of NPK ha<sup>-1</sup> + vermicompost @ 5 t ha<sup>-1</sup> + humic acid @ 0.2% + boron @ 1%. These findings lead towards beneficial and commercially feasible for the effective cultivation of tuberose (*Polianthes tuberosa* L.) cv. prajwal under open field conditions in the coastal ecosystems.

**Keywords:** Tuberose, organic manures, biostimulants, micronutrients, growth characteristics and flowering characteristics.

### Introduction

Tuberose (*Polianthus tuberosa* L.) is one of the most important tropical bulbous flowering plants cultivated for the production of long lasting flower spikes. It is popularly known as Rajanigandha. It belongs to the family Amaryllidaceae and it is native of Mexico. Commercial importance of tuberose is due to beauty of the flower, longer vase-life of spikes and aromatic oil extracted from its fragrant white flower (Alan *et al.*, 2007). The tuberose blooms throughout the year, florets are star-shaped, waxy and loosely arranged on spike that can reach up to 30 to 45 cm in length. There is high demand for tuberose concrete and absolute in international markets which fetch very good price. Flowers of the Single type (single row perianth) are commonly used as loose flowers, making garlands and essential oil etc, while the double varieties (more than two rows of perianth) are used as cut flowers, garden display and interior decoration (Anonymous, 2016).

Organic manures not only provide major nutrients but micronutrients as well to the growing plants. It increase the organic matter content and hence improve the physical properties of soils including water holding capacity in sandy soils and drain ability in clayey soils. Organic materials are the safer sources of plant nutrients which have no detrimental effect to crops and soil. Cow dung, farm yard manure, poultry manure, vermicompost and green manure are excellent sources of organic matter as well as primary plant nutrients (Pieters, 2005). Organic materials are the safer sources of plant nutrients which have no detrimental effect to crops and soil. Cow dung, farm yard manure, poultry manure, vermicompost and green manure are excellent

sources of organic matter as well as primary plant nutrients. Among all the animal manures, poultry manure has the highest amount of NPK content. Optimal storage conditions for chicken manure includes it being kept in a covered area and retaining its liquid, because a significant amount of nitrogen exists in the urine. Tuberose is a gross feeder and requires a large quantity of NPK both in the form of organic and inorganic fertilizers (Amarjeet *et al.*, 2000). A plant biostimulant is any substance or microorganism applied to plants with the aim to enhance nutrition efficiency, abiotic stress tolerance and/or crop quality traits, regardless of its nutrients content (Jardin, 2015). Micronutrients are essential for better growth of tuberose. The micronutrients are responsible in activating several enzymes (catalase, peroxidase, alcohol dehydrogenase, carbonic dehydrogenase, tryptophan synthase, etc.) and involve them self in chlorophyll synthesis and various physiological activities by which plant growth and development are encouraged (Kumar and Arora, 2000). The use of growth stimulants, organic manures and micronutrients has brought about a sort of revolution in the floriculture industry. Present research work was planned to investigate the best effect of organic manures, biostimulants and micronutrients of tuberose under open field conditions in the coastal ecosystems.

### Materials and Methods

The experiment was conducted in the agro-climatic conditions of Annamalai Nagar at floriculture yard, Department of Horticulture, Annamalai University, Chidambaram region features a hot summer climate with maximum temperature of 38°C and minimum of 17.5 °C with annual average rainfall of 1235 mm and consider good for

tuberoses cultivation. The corms of the single variety of tuberoses were sown after dipped in carbendazim fungicide 2 g litre<sup>-1</sup> of water for 15 minutes. The treatments were replicated three times during the month of February. The certain growth parameters including plant height, number of leaves plant<sup>-1</sup> and number of sideshoots plant<sup>-1</sup> were observed by using organic manures and foliar application of biostimulants and micronutrients. Randomized block design was used to evaluate the results statistically and LSD (least significant difference) at 5% were calculated according to the method described by Panse and Sukhatme (1978).

## Results and Discussion

### Growth characteristics

**Plant height (cm):** The data on plant height are shown in table 1. The plant height at different stages of growth differed significantly. Among various treatments, the maximum plant height was observed in T<sub>4</sub> (50 % RDF@ 100: 100: 100 kg NPK ha<sup>-1</sup> +vermicompost @ 5 t ha<sup>-1</sup> + humic acid @ 0.2 % + boron @ 1%) at 60 DAP (22.88 cm), 90 DAP (48.78 cm) and 120 DAP (64.84 cm). It was followed by T<sub>8</sub> (50% RDF @ 100: 100: 100 kg NPK ha<sup>-1</sup>+ poultry manure @ 10 t ha<sup>-1</sup> + seaweed extract @ 2000 g sq.m<sup>-1</sup> + zinc sulphate @ 0.5%), whereas the minimum was found to be in T<sub>13</sub> (control) at 60 DAP (15.52 cm), 90 DAP (39.62 cm) and 120 DAP (54.79 cm).

**Number of leaves plant<sup>-1</sup>:** The data with respect to number of leaves plant<sup>-1</sup> are presented in table 2. Among the treatments, plants treated with 50 % RDF@ 100: 100: 100 kg NPK ha<sup>-1</sup>+ vermicompost @ 5 t ha<sup>-1</sup> + humic acid @ 0.2 % + boron @ 1% (T<sub>4</sub>) showed the highest number of leaves plant<sup>-1</sup> at 60 DAP (28.71), 90 DAP (44.65) and 120 DAP (51.72), it was followed by T<sub>8</sub> (50% RDF @ 100: 100: 100 kg NPK ha<sup>-1</sup>+ poultry manure @ 10 t ha<sup>-1</sup> + seaweed extract @ 2000 g sq.m<sup>-1</sup> + zinc sulphate @ 0.5%). The minimum number of leaves plant<sup>-1</sup> at 60 DAP (21.42), 90 DAP (36.47) and 120 DAP (42.27) was found in T<sub>13</sub> (control).

**Number of side shoots plant<sup>-1</sup>:** The data with respect to the number of side shoots plant<sup>-1</sup> are furnished in table 3. The number of side shoots plant<sup>-1</sup> differed significantly by the effect of various treatments. The maximum number of side shoots plant<sup>-1</sup> at 60 DAP (7.21), 90 DAP (9.98) and 120 DAP (12.73) was observed in T<sub>4</sub> (50 % RDF@ 100: 100: 100 kg NPK ha<sup>-1</sup>+ vermicompost @ 5 t ha<sup>-1</sup> + humic acid @ 0.2 % + boron @ 1%) which was followed by T<sub>8</sub>. The minimum number of side shoots at 60 DAP (5.97), 90 DAP (6.24) and 120 DAP (7.98) were recorded in T<sub>13</sub> (control).

### Flowering Characteristics

**Spike length (cm):** The data relating to the effect of various integrated nutrient management treatments on length of spike are presented in table 4. Among the treatments, T<sub>4</sub> (50 % RDF@ 100: 100: 100 kg NPK ha<sup>-1</sup>+ vermicompost @ 5 t ha<sup>-1</sup>

+ humic acid @ 0.2 % + boron @ 1%) recorded the highest spike length (100.21 cm). It was followed by T<sub>8</sub> (98.98 cm). The lowest spike length (85.37 cm) was recorded in T<sub>13</sub> (control).

**Rachis length (cm):** The data on rachis length due to the effect of various treatments are furnished in table 4. Among the treatments, T<sub>4</sub> (50 % RDF@ 100: 100: 100 kg NPK ha<sup>-1</sup> + vermicompost @ 5 t ha<sup>-1</sup> + humic acid @ 0.2 % + boron @ 1%) recorded the maximum rachis length (37.26 cm) and the next best was with T<sub>8</sub> (50% RDF@ 100: 100: 100 kg NPK ha<sup>-1</sup> + poultry manure @ 10 t ha<sup>-1</sup> + seaweed extract @ 2000 g/sq.m + zinc sulphate @ 0.5%) which recorded 36.64 cm. The minimum rachis length (29.34 cm) was observed in T<sub>13</sub> (control).

**Number of floret spike<sup>-1</sup>:** The data pertaining to the effect of various treatments, on number of florets spike<sup>-1</sup> are furnished in table 5. Among the various treatments, T<sub>4</sub> (50 % RDF @ 100: 100: 100 kg NPK ha<sup>-1</sup> +vermicompost @ 5 t ha<sup>-1</sup> + humic acid @ 0.2 % + boron @ 1%) recorded more number of florets spike<sup>-1</sup> (53.23) and this was followed by T<sub>8</sub> (52.47 florets spike<sup>-1</sup>). The least number of florets (43.68 florets spike<sup>-1</sup>) were recorded in T<sub>13</sub> (control).

**Flower diameter (cm):** The data regarding floret diameter obtained by the impact of various treatments are presented in table 5. The highest value (5.16 cm) was observed in T<sub>4</sub> (50 % RDF @ 100: 100: 100 kg NPK ha<sup>-1</sup> + vermicompost @ 5 t ha<sup>-1</sup> + humic acid @ 0.2 % + boron @ 1%). The next highest value in flower diameter (5.08 cm) was recorded in T<sub>8</sub> (50% RDF @ 100: 100: 100 kg NPK ha<sup>-1</sup> + poultry manure @ 10 t ha<sup>-1</sup>+ seaweed extract @ 2%+ zinc sulphate @ 0.5%). The treatment T<sub>13</sub> (control) recorded the lowest value (3.30 cm).

Enhanced growth parameters observed in present experiment due to application of organic inputs to the soils can promote nutrients availability and plant uptake, increase crop yield, reduce inputs of chemical fertilizers and minimize environmental risks. The increased spike length might be due to rapid internode elongation as a result of increased in cell division and cell elongation in the intercalary meristem. Other flowering parameters observed in this study might be due to higher carbohydrate, other essential nutrients, plant growth regulators and enzymes deposition in flower cells by boron physiological role which resulted in production of good quality attractive flowers.

Hence these results were supported by the previous findings of Patil (2004) in tuberoses; Bhalla *et al.* (2006) in gladiolus; Suseela *et al.* (2016), Yathindra *et al.* (2016), Pal *et al.* (2017) in tuberoses, Misra (2001) in chrysanthemum, Sundaret *et al.* (2010) in *J. sambac* and Kumar *et al.* (2012) in gladiolus, Ganesh *et al.* (2014) for better growth and flowering parameters like plant height, number of leaves plant<sup>-1</sup> and number of sideshoots plant<sup>-1</sup>, spike length, rachis length, flower diameter and number of florets spike<sup>-1</sup>.

**Table 1 :** Effect of organic manures, biostimulants and micronutrients on plant height (cm) in tuberose (*Polianthes tuberosa* L.) cv. Prajwal

Treatments	Plant height (cm)		
	60 DAP	90 DAP	120 DAP
T <sub>1</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract @ 2% sq.m <sup>-1</sup> + boron @ 1%	21.65	47.21	63.10
T <sub>2</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	20.44	45.65	61.37
T <sub>3</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	18.73	43.46	58.94
T <sub>4</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	22.88	48.78	64.84
T <sub>5</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid + boron@ 1%	21.10	46.50	62.31
T <sub>6</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	19.89	44.94	60.58
T <sub>7</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	18.17	42.73	58.13
T <sub>8</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	22.31	48.05	64.03
T <sub>9</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	20.52	45.75	61.48
T <sub>10</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	19.32	44.22	59.78
T <sub>11</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	17.61	42.00	57.32
T <sub>12</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	22.23	47.95	63.92
T <sub>13</sub> - Control	15.52	39.62	54.79
<b>CD (0.05)</b>	<b>0.54</b>	<b>0.70</b>	<b>0.76</b>
<b>S.Ed</b>	<b>0.27</b>	<b>0.35</b>	<b>0.38</b>

Note: RDF- Recommended dosage of fertilizer @ 100:100:100 kg of NPK ha<sup>-1</sup>, humic acid, seaweed extract, boron and zinc as foliar application

**Table 2 :** Effect of organic manures, biostimulants and micronutrients on number of leaves plant<sup>-1</sup> in tuberose (*Polianthes tuberosa* L.) cv. Prajwal

Treatments	Number of leaves plant <sup>-1</sup>		
	60 DAP	90 DAP	120 DAP
T <sub>1</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract @ 2% sq.m <sup>-1</sup> + boron @ 1%	27.50	43.27	50.09
T <sub>2</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	26.30	41.90	48.48
T <sub>3</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	24.61	39.97	46.21
T <sub>4</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	28.71	44.65	51.72
T <sub>5</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid + boron@ 1%	26.95	42.64	49.35
T <sub>6</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	25.75	41.27	47.74
T <sub>7</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	24.05	39.33	45.46
T <sub>8</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	28.15	44.01	50.97
T <sub>9</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract @ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	26.37	41.98	48.58
T <sub>10</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	25.19	40.64	47.00
T <sub>11</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	23.49	38.64	44.71
T <sub>12</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	28.07	43.92	50.86
T <sub>13</sub> - Control	21.42	36.47	42.27
<b>CD (0.05)</b>	<b>0.53</b>	<b>0.60</b>	<b>0.72</b>
<b>S.Ed</b>	<b>0.27</b>	<b>0.30</b>	<b>0.36</b>

Note: RDF- Recommended dosage of fertilizer @ 100:100:100 kg of NPK ha<sup>-1</sup>, humic acid, seaweed extract, boron and zinc as foliar application

**Table 3 :** Effect of organic manures, biostimulants and micronutrients on number of side shoots plant<sup>-1</sup> in tuberose (*Polianthes tuberosa* L.) cv. Prajwal

Treatments	Number of sideshoots plant <sup>-1</sup>		
	60 DAP	90 DAP	120 DAP
T <sub>1</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract @ 2% sq.m <sup>-1</sup> + boron @ 1%	6.97	9.29	12.01
T <sub>2</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	6.73	8.81	11.29
T <sub>3</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	6.40	8.12	10.29
T <sub>4</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	7.21	9.98	12.73
T <sub>5</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid + boron@ 1%	6.86	9.07	11.68
T <sub>6</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	6.63	8.58	10.96
T <sub>7</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	6.29	7.90	9.95
T <sub>8</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	7.10	9.55	12.40
T <sub>9</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	6.75	8.84	11.34
T <sub>10</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	6.52	8.36	10.63
T <sub>11</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	6.18	7.68	9.61
T <sub>12</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	7.08	9.52	12.35
T <sub>13</sub> - Control	5.97	6.24	7.98
<b>CD (0.05)</b>	<b>0.10</b>	<b>0.20</b>	<b>0.30</b>
<b>S.Ed</b>	<b>0.05</b>	<b>0.10</b>	<b>0.15</b>

Note: RDF- Recommended dosage of fertilizer @ 100:100:100 kg of NPK ha<sup>-1</sup>, humic acid, seaweed extract, boron and zinc as foliar application.

**Table 4 :** Effect of organic manures, biostimulants and micronutrients on spike length (cm) and rachis length(cm) in tuberose (*Polianthes tuberosa* L.) cv. Prajwal

Treatments	Spike length(cm)	Rachis length (cm)
T <sub>1</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract @ 2% sq.m <sup>-1</sup> + boron @ 1%	97.54	35.93
T <sub>2</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	94.90	34.61
T <sub>3</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	91.19	32.75
T <sub>4</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	100.21	37.26
T <sub>5</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid + boron@ 1%	96.33	35.32
T <sub>6</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	93.69	34.00
T <sub>7</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	89.96	32.13
T <sub>8</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	98.98	36.64
T <sub>9</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	95.06	34.69
T <sub>10</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	92.47	33.39
T <sub>11</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	88.73	31.51
T <sub>12</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	98.80	36.56
T <sub>13</sub> - Control	85.37	29.34
<b>CD (0.05)</b>	<b>1.18</b>	<b>0.58</b>
<b>S.Ed</b>	<b>0.59</b>	<b>0.29</b>

Note: RDF- Recommended dosage of fertilizer @ 100:100:100 kg of NPK ha<sup>-1</sup>, humic acid, seaweed extract, boron and zinc as foliar application

**Table 5 :** Effect of organic manures, biostimulants and micronutrients on number of floret spike<sup>-1</sup> and flower diameter (cm) in tuberose (*Polianthes tuberosa* L.) cv. Prajwal

Treatments	Flower diameter (cm)	Number of floret spike <sup>-1</sup>
T <sub>1</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract @ 2% sq.m <sup>-1</sup> + boron @ 1%	4.99	51.58
T <sub>2</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	4.82	49.95
T <sub>3</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + boron@ 1%	4.59	47.66
T <sub>4</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	5.16	53.23
T <sub>5</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid + boron@ 1%	4.91	50.84
T <sub>6</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + boron@ 1%	4.75	49.20
T <sub>7</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	4.51	46.90
T <sub>8</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	5.08	52.47
T <sub>9</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + seaweed extract@ 2% sq.m <sup>-1</sup> + zinc sulphate@ 0.5%	4.83	50.05
T <sub>10</sub> - 50% RDF + vermicompost@ 5 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	4.67	48.45
T <sub>11</sub> - 50% RDF + poultry manure@ 10 t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	4.43	46.14
T <sub>12</sub> - 50% RDF + neem cake@ 1t ha <sup>-1</sup> + humic acid@ 0.2% + zinc sulphate@ 0.5%	5.07	52.36
T <sub>13</sub> - Control	3.30	43.68
<b>CD (0.05)</b>	<b>0.05</b>	<b>0.70</b>
<b>S.Ed</b>	<b>0.02</b>	<b>0.35</b>

Note: RDF- Recommended dosage of fertilizer @ 100:100:100 kg of NPK ha<sup>-1</sup>, humic acid, seaweed extract, boron and zinc as foliar application

### Conclusion

In light of the above discussions, it could be concluded that foliar application of 50 % RDF @ 50: 50: 50 kg NPK ha<sup>-1</sup> + vermicompost @ 5 t ha<sup>-1</sup> along with foliar application of humic acid @ 0.2 % + boron @ 1% at 60 days interval from 60 DAP onwards was found better in increasing growth and flowering yield of tuberose cv. Prajwal.

### References

- Alan, O.; Gunen, Y.; Ceylan, S. and Gunen, E. (2007). Effect of nitrogen applications on flower yield, some quality characteristics and leaf mineral content in tuberose (*Polianthes tuberosa* L.), Ege Tarimsal Arastirma Enstitusu Mudurlugu, Izmir, Turkey: Aegean Agriculture Research Ins. Direc., 17(1): 43-57.
- Amarjeet, S.; Godara, N.R.; Ashok, K.; Singh, A. and Kumar, A. (2000). Effect of NPK on flowering and flower quality of tuberose (*Polianthes tuberosa* L.) cv. Single. Haryana Agriculture University Journal Research, 26(1): 43-49.
- Anonymos (2016). Horticulture: flower crops: Tuberose <http://www.Agritech.Tnau.ac.in/horticulture>.
- Bhalla, R.; Kanwar, P.; Dhiman, S.R. and Jain, R. (2006). Effect of biofertilizers and biostimulants on growth and flowering in gladiolus. Journal of Ornamental Horticulture, 9(4): 248-252.
- Ganesh, S.; Soorianathasundaram, K.; Kannan, M. and Jeyakumar, P. (2014). Influence of plant growth regulators and micronutrients on growth, floral characters and yield of tuberose (*Polianthes tuberosa* L.) cv. Prajwal. J. Ornamental Hort., 15(1&2): 696-700.
- Jitendra, K.; Amin, M. and Singh, P.V. (2003). Effect of Mn and Zn spray on carnation. J. Ornamental Hort., 6(1): 83.
- Kumar, P. and Arora, J.S. (2000). Effects of Micronutrients on Gladiolus. J. Orn. Hort. (New Series), 3: 91-93.
- Misra, H.P. (2001). Response of chrysanthemum to zinc and boron on growth, yield and quality of flowers. Scientific Hort., 7: 201-208.
- Pal, A.K.; Ragupathi, B.; Nellipalli, V.K. and Karim, K.B. (2017). Effects of biostimulants on growth and floral attributes of Tuberose (*Polianthes tuberosa* L.) cv. Prajwal. Int. J. Curr. Microbiol. App. Sci., 6(6): 66-69.
- Panse, V.G. and Sukhatme, P.V. (1978). Statistical methods for Agricultural workers, Indian Council of Agrl. Res., New Delhi, 3<sup>rd</sup>edn.
- Patil, S.R.; Reddy, B.S. and Prasanth, J.M. (2004). Effect of organic, inorganic and insitu, vermicompost on chlorophyll content and flower yield of *Jasminum sambac* Ait. J. Orn. Hort., 7(3-4): 164-167.
- Pieters, A.J. (2005). Green manuring: Principles and practices. Agrobios, Jodhpur, India.
- Suseela, T.; Chandrasekar, R.V.; VijayaBhaskar, D.R. Salomi Suneetha and Krishna, K.U. (2016). Effect of organic manure inorganic and micronutrient on vegetative and floral characters of tuberose (*Polianthes tuberosa* L.) cv- Suvasini, Int. J. of Scientific and Research Pulication, 6(1): 170 – 173.
- Yathindra, H.A.; Krishna, M.R.; Rajesh, A.M. and Harshavardhan, M. (2016). Effect of integrated nutrient management on growth parameters of Bird of Paradise (*Strelitzia reginae* L.). The Bioscan, 11(1): 565-568.