



Plant Archives

Journal homepage: <http://www.plantarchives.org>
 DOI Url : <https://doi.org/10.51470/PLANTARCHIVES.2022.v22.no1.014>

ESTIMATION OF ANTIOXIDATIVE ENZYMES IN THE TESTES AND LIVER OF THE *HETEROPNEUSTES FOSSILIS* EXPOSED TO PENTACHLOROPHENOL

Sunanda Dubey^{1*}, Surya Prakash Verma² and Shashikant Dwivedi³

^{1,2}Department of Zoology, Udai Pratap College, Varanasi-221002 India,

³Department of Physics, Udai Pratap College, Varanasi-221002, India.

*Corresponding author: dubey.sunanda@gmail.com

(Date of Receipt : 08-08-2021; Date of Acceptance : 25-11-2021)

ABSTRACT

Oxidative stress has become an area of significant interest for toxicological studies. Estimation of antioxidative enzymes such as Catalase (CAT), Superoxide dismutase (SOD), Lipid peroxidase (LPO) and Glutathion peroxidase (GPx) have been measured in the fish exposed to 1/10th LC₅₀ (32µg/l) PCP for 14 and 28 days. SOD activity and CAT activity in testes and liver were found to be decreased as compared to control groups, whereas levels of Malondialdehyde (MDA) and GPx in testicular and hepatic tissue were significantly elevated as compared to control groups of catfish. Results showed oxidative stress alteration due to PCP exposure. Oxidative damage is produced in response to the exposure to environmental contaminants, by generating free radicals and altering antioxidant enzyme systems. The results of the present investigation indicate that exposure of PCP induces significant changes in the enzymatic profiles in *Heteropneustis fossilis*. The presence of such level of PCP in the natural environment is dangerous to the ecosystem and will definitely affect the survival of fish.

Keywords : Antioxidative enzymes, Catfish, *H. fossilis*, Oxidative stress.

Introduction

Extensive use of pesticides finds their way to aquatic environment and becomes a serious problem and affects the health and survival of non-target organism. Oxidative stress is more often used as a biomarker to study the effects of environmental pollution in aquatic environments. Compounds that can inhibit or prevent the oxidation of oxidizable materials by scavenging free radicals and reducing oxidative stress are defined as Antioxidants. Many pollutants may exert toxicity related to oxidative stress. The level of antioxidant enzymes in organism have been extensively taken as an early warning indicator of pollution (Lin *et al.*, 2001). Now days, most environmental problems are attributed to the production and release of toxic chemicals which are capable of interacting with the environment and disrupting the ecosystem. The pentachlorophenol (PCP) known to be a widely distributed persistent environmental contaminant (Heudorf *et al.*, 2000; Ge *et al.*, 2007; Geyer *et al.*, 1987; WHO, 1987; Muir and Eduljee, 1999; OEHHA, 1997). The PCP exposure has been associated with many adverse effects on health and the eco environment (WHO, 1987; Verma and Dubey, 2019, 2021). Based on inadequate evidence in humans but sufficient evidence in animal studies, the International Agency for Research on Cancer classified PCP as a group 2B carcinogen (IARC, 1991). Potential thyroid disrupting effects of PCP have also been revealed (Dallaire *et al.*, 2009; Van den Berg, 1990).

The global restriction on PCP production and use has resulted in reduced PCP contamination in the environment in different world regions (Muir and Eduljee, 1999). In China the primary use of PCP and its sodium salt (Na-PCP) is to kill Schistosome intermediate host snails (Ge *et al.*, 2007; CESE, 2004), wood preservation and other uses account for the secondary (Zhou *et al.*, 2005). The increased PCP use could be resulted in more environmental contamination and potential health risks. In addition to the health risks caused by PCP itself, the impurities of commercial Na-PCP, polychlorinated dibenzo-p-dioxins and dibenzofurans (PCDD/Fs), can be released into the environment with the use of Na-PCP, this compounds the health risks caused by PCP (Bao *et al.*, 1995; Zheng *et al.*, 2008). Analysis of biomarkers in aquatic organisms particularly in fish is a validated approach for early warning of chemical exposure (Lin *et al.*, 2001; Van der Oost *et al.*, 2003; Osman *et al.*, 2010). It is suggested that during stress conditions fish are capable to change and adapt their metabolic functions (Malarvizhi *et al.*, 2012). The inhibition or induction of the enzymes such as glutamate oxaloacetate transaminases (GOT), glutamate pyruvate transaminase (GPT), lactate dehydrogenase (LDH), acid phosphatase (ACP), alkaline phosphates (ALP) etc., can be used to indicate tissue damage (Nemcsok and Boross, 1982; Webb *et al.*, 2005). Enzymes play important roles in protecting the cell against the potentially toxic effects of the environment (Kuthan *et al.*,

1986), superoxide dismutase catalyzes the dismutation of the superoxide ion (O_2^-) to hydrogen peroxide and oxygen molecule during oxidative energy processes. GPT, GOT and LDH functions as link enzymes between the protein and carbohydrate metabolism, and also serve as an indicator of chemical stress. Moreover, pesticides enhanced the production of reactive oxygen species (ROS) in aquatic organisms which may lead to impairment of many cellular functions (Uner *et al.*, 2006; Monteiro *et al.*, 2009; Modesto and Martinez, 2010; De Menezes *et al.*, 2011). Antioxidant systems can prevent or counterbalance the deleterious effect of free radicals (Lushchak, 2011; Singh *et al.*, 2011). Fish are reported to possess both mitochondrial and cytosolic superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx) and non-enzymatic (reduced glutathione and vitamin E) antioxidants as in mammals (Droge, 2002) for detoxification of free radicals (Lushchak, 2011). It is stated that the level of enzyme SOD and CAT can be used as markers of pollutant induced oxidative stress in aquatic organisms (Borkovi *et al.*, 2005). These antioxidants are essential to maintain the redox status of fish cells, likewise, alterations in Lipid peroxidation (LPO) concentration are also used to express severe oxidative damage and are used as a biomarker of effect (Van der Oost *et al.*, 2003; Lackner, 1998). Malondialdehyde (MDA) is a naturally occurring product of LPO which acts as its indicator (Charissou *et al.*, 2004; Pampanin *et al.*, 2005). Fish are obviously subjected to exposure to different pollutants often present in the aquatic environment (Velkova-Jordanoska *et al.*, 2008). Fish has been used as indicator species for monitoring of pollution in the aquatic environment. *H. fossilis*, a catfish, is used for this study to determine the health status by estimating the antioxidative enzymes in the testes and liver of the fish exposed to PCP.

Material and Methods

Experimental setup

Pentachlorophenol (Crystalline, 99% pure) was purchased from Acros organics (Geel, Belgium). Other chemicals used were of analytical grade and purchased locally. Pentachlorophenol was dissolved in ethanol and then diluted with water to obtain the required concentrations $32\mu\text{g/l}$ ($1/10^{\text{th}}$ of LC_{50}).

Estimation of LC_{50}

The freshwater male catfish, *H. fossilis* were purchased from the local market in Varanasi and used for the experiment. Estimation of LC_{50} for PCP, the protocol of Secretaria Estadual Do Meio Ambiente (SEMA, 1988) was followed. The LC_{50} value for fish was $320\mu\text{g/l}$ and for further experiments $1/10^{\text{th}}$ LC_{50} ($32\mu\text{g/l}$) dose were selected (Pandey and Dubey, 2015). Mortality was observed to be 3 in number. They were maintained in the laboratory under normal photo period (13.0L: 11.0D) and temperature ($25\pm 2^{\circ}\text{C}$) until use for experiments. The fish were fed with boiled egg daily *ad libitum*. The experiments were performed in accordance with the guidelines for experimentation in animals and all care was taken to prevent cruelty of any kind.

The acclimatized adult male fish classified into two groups (15 fish per each): first group control, second group PCP treated (for 14 days and 28 days with $32\mu\text{g/l/day}$). In the present study, the amount of PCP exposures was $32\mu\text{g/l}$ and the exposure concentrations were determined by performing

LC_{50} experiment. The conditions of the experiment were as that of acclimatization with changing all the containers water and concentrations of PCP every day. Male catfish *H. fossilis* (30-45g) were purchased from local fish market in the Resting phase for the experiment.

Measurement of antioxidant status in liver and testes

Estimation of Catalase (CAT) activity

CAT activity was assayed by the method of Chance and Maehly (Chance and Maehly, 1955). The liver and testes were homogenized (10%) in 50 mM phosphate buffer, pH 7.0, and centrifuged at 16,000g for 45 min. The supernatant was used for the enzyme assay. The reaction mixture contained 2 ml of phosphate buffer (pH 7.0), 0.45 ml H_2O_2 , and 0.025 ml of enzyme source. The enzyme activity was expressed as micromoles of H_2O_2 metabolized/milligram protein/minute, at 250 nm of absorbance.

Estimation of Super oxide dismutase (SOD) activity

SOD activity was measured as the inhibition of photo reduction of nitrobluetetrazolium (NBT) by the enzyme as per the method of Beauchamp and Fridovich (Beauchamp and Fridovich, 1971). The tissues were homogenized in (10% w/v) potassium phosphate buffer (pH 7.5) containing 1% polyvinyl pyrrolidone and centrifuged at 16,000g for 15 min. The supernatant was used as the enzyme source. The total reaction mixture consisted of 100 mM phosphate buffer, 10mM EDTA, 130 mM methionine, 750 mM NBT, 60 mM riboflavin, and enzyme source. The reaction was initiated by the addition of riboflavin, the samples were placed under fluorescence for 30 min, and the resulting color was read at 560 nm against a reagent blank kept in a dark place. The activity was expressed as units/milligram protein.

Estimation of Glutathione peroxidase (GPx) activity

GPx activity was assayed by the method of Paglia and Valentine (Paglia and Valentine, 1967) with modifications according to Lawrence and Burke (Lawrence and Burke, 1976). The reaction mixture contained 50 mM potassium phosphate buffer (pH 8.3), 1 mM EDTA, 1mM sodium azide, 0.2 mM NADPH, and 1 U/ml glutathione reductase. The reaction was initiated with the addition of 1.5 m Mcumenehydroperoxide. The enzyme activity was estimated from the rate of oxidation of NADPH ($E_{340}=6200\text{M}^{-1}\text{cm}^{-1}$) and the enzyme activity was expressed as mmol/minute/milligram protein.

Lipid peroxidation (LPO) concentration

For the evaluation of LP concentration we measured malondialdehyde (MDA) by the TBARS method at 535 nm (Buege and Aust, 1978). 2 ml of the reaction mixture [thiobarbituric acid (0.375%), trichloroacetic acid (15%) and hydrochloric acid (0.25 N)] were mixed in 1:1:1 and were added to 1 ml of the heat denatured supernatant. TBARS levels were estimated at 535 nm using MDA as standard. Total protein concentration was measured on the supernatants (10,000g during 20 min at 4°C) of the homogenized tissues (Stoscheck, 1990). The concentration of LP compounds was expressed as n moles of MDA formed per mg protein.

Statistical analysis

Data were analyzed through a one-way analysis of variance (ANOVA) followed by post hoc test, Tukey's

multiple range test ($P < 0.05$). Data were expressed as mean \pm standard error mean (SEM).




Fig. 1 : Catalase (CAT) level in the liver and testes of a freshwater catfish, *H. fossilis* exposed to PCP for 14 and 28 days. (Asterisks shows significant difference from Control at $P < 0.05$; ANOVA following Tukey's post hoc Test)




Fig. 2 : Super Oxide Dismutase (SOD) level in the liver and testes of a freshwater catfish, *H. fossilis* exposed to PCP for 14 and 28 days. (Asterisks shows significant difference from Control at $P < 0.05$; ANOVA following Tukey's post hoc Test)




Fig. 3 : Glutathione Peroxidase (GPx) level in the liver and testes of a freshwater catfish, *H. fossilis* exposed to PCP for 14 and 28 days. (Asterisks shows significant difference from Control at $P < 0.05$; ANOVA following Tukey's post hoc Test)




Fig. 4 : Lipid Peroxidase (LPO) level in the liver and testes of a freshwater catfish, *H. fossilis* exposed to PCP for 14 and 28 days. (Asterisks shows significant difference from Control at $P < 0.05$; ANOVA following Tukey's post hoc Test)

Results

SOD (0.48; 0.34 units/milligram protein) activity was found to be decreased as compared to control groups (1.34; 0.86 units/milligram protein) and CAT activity was also observed to be lower in testes and liver (7.12; 10.07 micromoles of H_2O_2 metabolized/milligram protein/minute) than in the control groups (25.61; 23.08 micromoles of H_2O_2 metabolized/milligram protein/minute) respectively.

The findings showed elevated level of MDA in testicular (4.09 nmoles of MDA per gram of wet mass) and hepatic tissue (3.13 nmoles of MDA per gram of wet mass) than in control testes (0.86 nmoles of MDA per gram of wet mass) and liver (0.45 nmoles of MDA per gram of wet mass) of fish signifying free radical induced cell injury due to oxidative stress. GPx was also found to be significantly high in testicular (3.86 mmol/minute/milligram protein) and hepatic tissue (3.52 mmol/minute/milligram protein) than in control testes (1.34 mmol/minute/milligram protein) and liver (1.19 mmol/minute/milligram protein) respectively.

Discussion

Various industries (pulp, rubber, glass, sugar, tannery, steel plants, chemicals, plastics etc.) alongside the banks of river discharge the waste directly into water body. Studies on pollutant-induced toxicity in fish have revealed bioaccumulation of heavy metals in different tissues (Sen *et al.*, 2011; Fatima and Usmani, 2013) thereby, affecting the cellular architecture and physiology. Hence, study of fish as affected animal provides an insight into the potential toxic effects of perilous aquatic environments (Tyor and Pahwa, 2017). The exposure to environmental contaminants may produce oxidative damage by generating free radicals and/or altering antioxidant enzyme systems (Huang *et al.*, 2007). Free radical (molecule with one or more unpaired electrons in a single orbit) produced as a result of ATP production by mitochondria become a part of the propagative chain reaction whereby they combine with other radicals to form other more damaging species, unless the chain is terminated by antioxidants to form a nontoxic species (Halliwell, 1994). The free radical immediately react with lipids in the membranes of cell and organelles, proteins and DNA eventually forming damaging products such as lipid peroxides and other lipid adducts. The indicator of lipid peroxidation includes measurement of TBARS content (Wheatley, 2000) which provides relative measure of the potential for pollutants to cause oxidative injury

(Vlahogianni *et al.*, 2007). During normal physiological status anti-oxidant defense systems provides compensatory response to oxidative damage by removing reactive oxygen species (Livingstone, 2001) and thus, have been considered as biomarkers of contaminant, its induction reflects a protective response to pollutants (Borkovic *et al.*, 2005). An increase in the LPO level was observed after fluoride exposure in the liver and ovary of fishes (Basha and Sujitha, 2012). Increase in MDA was an indicator of enhanced oxidative stress (Buyukokuroglu *et al.*, 2002). Significant increase in MDA content was observed in Dichlorvos exposed *H. fossilis* (Vadhva and Hasan, 1986). Increased MDA level was also observed in the study carried out by scientist (Hai *et al.*, 1997) in *Cyprinus carpio* and *Ictalurus nebulosus* following the exposures to dichlorvos. SOD has been known as the first defense line of cell against ROS and could protect against superoxide induced oxidative damage (Fridovich, 1989) by catalyzing the conversion of the superoxide anion radical to molecular oxygen and hydrogen-peroxide (H_2O_2) while, CAT is involved in mitigating the toxic effects of H_2O_2 by decomposing it into water and oxygen. Inhibition of catalase activity may be due to overproduction of superoxide anion radical (Pandey *et al.*, 2003) or due to presence of nitrites (Arrillo and Melodia, 1991).

In the present investigation tissue specific responses in the activities of biomarker antioxidant enzymes such as SOD which detoxifies toxic superoxide anion radicals, catalase known to be primary antioxidant defense component, GPx and LPO were observed for assessing the PCP toxicity.

The findings of present study revealed significant decrease in the activities of SOD, CAT enzymes in liver and testes of fish when compared to control. Thus, the decrease in both the enzymes may be suggestive of the reason that their activity might be inhibited by the pollutants. The breakdown of H_2O_2 to water and oxygen is done by CAT, thereby, protecting the cell from the damaging action of H_2O_2 and the hydroxyl radical. SOD and CAT level decreased significantly in testes and liver when exposed to $32\mu g/l$ concentration of PCP. The PCP exposure increased the GPx and LPO activity in testes and liver of fish compared to control group. The elevation of antioxidant activities in organism might be indicative of the adaptive response of organisms to counteract the oxidative effect of reactive oxygen species generated by PCP exposure.

Reports discussed that GPx catalyzes the reaction of hydro peroxides with reduced GSH to form glutathione disulfide and the reduction product of the hydro peroxide and they further added that the decrease in activity of aminotransferases in the liver of fish *Oreochromis mossambicus* exposed to organo-phosphorus pesticide may be due to liver damage (Rao, 2006). In a recent study the decreased GOT and GPT in gill, liver and muscle of fish *Cyprinus carpio* after carbamazepine exposure suggested that detoxification mechanism may not be sufficiently effective to prevent the animal from toxicity and effect of drug on the system (Malarvizhi *et al.*, 2012). Furthermore, in some research it is suggested that the decrease in the activities of the metabolic enzymes during organophosphorus pesticides exposure may be due to the direct action of these pesticides on enzymes (Tripathi and Shasmal, 2011). Both mammalian and piscine systems shows general mechanisms of oxidative toxicity which appear more or less similar with comparable lesions

and its responses, serve as markers of oxidative stress (Vutukuru *et al.*, 2006). The oxidative stress caused by pesticides in aquatic organisms may lead to ROS production and alterations in antioxidant enzymes (Livingstone, 2001). The excessive production of ROS production and their damaging effects can be minimized by the cellular antioxidant systems (Sureda *et al.*, 2004; Dorts *et al.*, 2012). Scientists also advocated that organisms under stress conditions utilizes the antioxidant enzymes to adapt to environmental stress and altered activities depend on the dose, species and route of exposure (Gravato *et al.*, 2006). The enzymes such as catalase and SOD play a major role in counteracting the ROS produced during bioactivation of Xenobiotics and the induction of CAT/SOD system might be the foremost defense mechanism against ROS (Nwani *et al.*, 2010). Moreover, it is stated in some reports that SOD and CAT are highly sensitive enzymes respond quickly to protect organisms from prevailing oxidative stress (Rao *et al.*, 2006; Dewez *et al.*, 2005). SOD is the first enzyme that responds to oxidative stress during any stress condition in animals (McCord and Fridovich, 1969; Winston and Di Giulio, 1991). The elevation of antioxidant activities in organism generally indicates adaptive responses of organisms to counteract the oxidative effect of generated reactive oxygen species (Hegazi *et al.*, 2010).

Decrease in catalase activity observed in the liver and testes of fish in exposed group may be due to its adaptive response to H_2O_2 produced by SOD activity, since, catalase is responsible for its detoxification to water (Nwani *et al.*, 2010; Elia *et al.*, 2002). The variation in the activity of antioxidant enzymes may as indicators of pollutant mediated oxidative stress (Sayeed *et al.*, 2003). MDA is the final product of lipid peroxidation and their concentration in animals can be taken as direct evidence of toxic process caused by free radicals (Sieja and Talerczyk, 2004). The elevation of LPO content in and liver indicates an increase in free radicals due to toxicity or metabolism (Freeman and Crapo, 1981; Gultekin and Akdogan, 2000; Ruas *et al.*, 2008). The results of the present investigation indicate that exposure of PCP induces significant changes in the enzymatic profiles in *H. fossilis*. The PCP exposure to the fish indicated the oxidative stress in the body by generating imbalance between reactive oxygen species (ROS) and antioxidants. The presence of such level of PCP in the natural environment is dangerous to the ecosystem and will definitely affect the survival of fish. We are further interested in studying the assessment of residual amount of this pesticide in different body tissues of fish for further understanding.

Acknowledgement

The authors thank Professor R. Chaube, Department of Zoology, BHU, Varanasi for extending her valuable help in performing enzymatic analysis in her lab.

References

- Arrillo, A. and Melodia, F. (1991). Nitrite oxidation in *Eisema foetida* (Savigny): ecological implications. *Funct Ecol*, 5: 629–634.
- Bao, Z.C.; Wang, K.O.; Kang, J.X. and Zhao, L.W. (1995). Analysis of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in pentachlorophenol and sodium pentachlorophenol. *Environ Chem*, 14: 317–21.

- Basha, P.M. and Sujitha, N.S. (2012). Combined influence of intermittent exercise and temperature stress on the modulation of fluoride toxicity. *Biol Trace Elem Res*, 148: 69–75.
- Beauchamp, C. and Fridovich, I. (1971). Superoxide dismutase: Improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry*, 44: 276–287.
- Borkovi, S.S.; Pavlovic, S.Z.; Kovacevic, T.B.; Stajn, A.S.; Petrovic, V.M. and Saicic, Z.S. (2005). Antioxidant defense enzyme activities in hepatopancreas, gills and muscle of spiny cheek crayfish, *Orconectes limosus* from the river Danube. *Comp Biochem Physiol C*, 147: 122–128.
- Borkovic, S.S.; Saponjic, J.S.; Pavlovic, S.Z.; Blagojevic, D.P.; Milosevic, S.M.; Kovacevic, T.B.; Radojicic, R.M.; Spasic, M.B.; Zikic, R.V. and Saicic, Z.S. (2005). The activity of antioxidant defence enzymes in the mussel *Mytilus galloprovincialis* from the Adriatic Sea. *Comp Biochem Physiol*, 141: 366–374.
- Buege, J.A. and Aust, S.D. (1978). Microsomal lipid peroxidation. *Methods in Enzymology*, 52: 302–310.
- Buyukokuroglu, M.E.; Taysi, S.; Polat, F. and Gocer, F. (2002). Mechanism of the beneficial effects of dantrolene sodium on ethanol-induced acute gastric mucosal injury in rats. *Pharmacol Res*, 45: 421–425.
- CESE (College of Environmental Sciences and Engineering, Peking University) (2004). The survey report of pesticide POPs production industry. Beijing: Convention Implementation Office, State Environment Protection Administration, (in Chinese).
- Chance, B. and Maehly, A.C. (1955). Assay of Catalase and Peroxidases. *Methods in Enzymology*, 2: 764–775.
- Charissou, A.M.; Cossu-Leguille, C. and Vasseur, P. (2004). Relationship between two oxidative stress biomarkers, malondialdehyde and 8-oxo-7,8-dihydro-20-deoxyguanosine, in the freshwater bivalve *Unio tumidus*. *Sci Total Environ*, 322: 109–122.
- Dallaire, R.; Muckle, G.; Dewailly, É.; Jacobson, S.W.; Jacobson, J.L.; Sandanger, T.M.; Sandau, C.D. and Ayotte, P. (2009). Thyroid hormone levels of pregnant Inuit women and their infants exposed to environmental contaminants. *Environ Health Perspect*, 117: 1014–1020.
- De Menezes, C.C.; Loro, V.L.; da Fonseca, M.B.; Cattaneo, R.; Pretto, A.; dos Santos Miron, D.; and Santi, A. (2011). Oxidative parameters of *Rhamdia quelen* in response to commercial herbicide containing clomazone and recovery pattern. *Pestic. Biochem. Physiol*, 100: 145–150.
- Dewez, D.; Geoffroy, L.; Vernet, G. and Popovic, R. (2005). Determination of photosynthetic and enzymatic biomarkers sensitivity used to evaluate toxic effects of copper and fludioxonil in alga, *Scenedesmus obliquus*. *Aquat Toxicol*, 74: 150–159.
- Dorts, J.; Bauwin, A.; Kestemont, P.; Jolly, S.; Sanchez, W. and Silvestre, F. (2012). Proteasome and antioxidant responses in *Cottus gobio* during a combined exposure to heat stress and cadmium. *Comp Biochem Physiol C*, 155: 318–324.
- Droge, W. (2002). Free radicals in the physiological control of cell function. *Physiol Rev*, 82: 47–95.
- Elia, A.C.; Waller, W.T. and Norton, S.J. (2002). Biochemical responses of bluegill sunfish (*Lepomis macrochirus*, Rafinesque) to atrazine induced oxidative stress. *Bull Environ Contam Toxicol*, 68: 809–816.
- Fatima, M. and Usmani, N. (2013). Histopathology and bioaccumulation of heavy metals (Cr, Ni and Pb) in fish (*Channa striatus* and *Heteropneustes fossilis*) tissue: a study for toxicity and ecological impacts. *Pak J Biol Sci*, 16: 412–420.
- Freeman, B.A. and Crapo, J.D. (1981). Hyperoxia increases oxygen radical production in rat lung and lung mitochondria. *J Biol Chem*, 256: 10986–10992.
- Fridovich, I. (1989). Superoxide dismutases. An adaptation to a paramagnetic gas. *J Biol Chem*, 264: 7761–7764.
- Ge, J.; Pan, J.; Fei, Z.; Wu, G. and Giesy, J.P. (2007). Concentrations of pentachlorophenol (PCP) in fish and shrimp in Jiangsu Province, China. *Chemosphere*, 69(1): 164–169.
- Geyer, H.J.; Scheunert, I. and Korte, F. (1987). Distribution and bio-concentration potential of the environmental chemical pentachlorophenol (PCP) in different tissues of humans. *Chemosphere*, 16: 887–99.
- Gravato, C.; Teles, M.; Oliveira, M. and Santos, M.A. (2006). Oxidative stress, liver biotransformation and genotoxic effects induced by copper in *Anguilla anguilla* L. – the influence of pre-exposure to b naphthoflavone. *Chemosphere*, 65: 1821–1830.
- Gultekin, F.M. and Akdogan, M. (2000). The effect of organophosphate insecticide chlorpyrifos-ethyl on lipid peroxidation and antioxidant enzymes (in vitro). *Arch Toxicol*, 74: 533–538.
- Hai, D.Q.; Varga, S.I. and Matkovic, B. (1997). Organophosphate effects of antioxidant system of carp (*Cyprinus carpio*) and catfish (*Ictalurus nebulosus*). *CompBiochemPhysiol*, 117: 83–8.
- Halliwell, B. (1994). Free radicals, antioxidants, and human disease: curiosity, cause, or consequence? *Lancet*, 344: 721–724.
- Hegazi, M.M.; Attia, Z.I. and Ashour, O.A. (2010). Oxidative stress and antioxidant enzymes in liver and white muscle of *Nile tilapia* in chronic ammonia exposure. *Aquat Toxicol*, 99: 118–125.
- Heudorf, U.; Letzel, S.; Peters, M. and Anger, J. (2000). PCP in the blood plasma: Current exposure of the population in Germany based on data obtained in 1998. *Int J HygEnviron Health*, 203: 135–9.
- Huang, G.J.; Chen, H.J.; Chang, Y.S.; Sheu, M.J. and Lin, Y.H. (2007). Recombinant sporamin and its synthesized peptides with antioxidant activities in vitro. *Bot Stud*, 48: 133–140.
- IARC (1991). Occupational exposures in insecticide application and some pesticides. Lyon, International Agency for Research on Cancer, pp. 371–402 (IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 53).
- Kuthan, H.; Haussmann, H.J. and Werringlover, J. (1986). A spectrophotometric assay for superoxide dismutase activities in crude tissue fractions. *Biochem J*, 237: 175–180.
- Lackner, R. (1998). Oxidative stress. In: Fish by Environmental Pollutants, Fish Ecotoxicology, (Ed. T Braunbeck, DE Hinton and B Streit, Berlin, Germany) 203–24.
- Lawrence, R.A. and Burke, R.F. (1976). Glutathione peroxidase activity in selenium-deficient rat liver.

- Biochemical and Biophysical Research Communications*, 71: 952–958.
- Lin, C.T.; Lee, T.L.; Duan, K.J. and Su, J.C. (2001). Purification and Characterization of Black Porgy Muscle Cu/Zn Superoxide Dismutase. *Zoological Studies*, 40: 84-90.
- Livingstone, D.R. (2001). Contaminant stimulated reactive oxygen species production and oxidative damage in aquatic organisms. *Mar Pollut Bull*, 42: 656–666.
- Lushchak, V.I. (2011). Environmentally induced oxidative stress in aquatic animals. *Aquat Toxicol*, 101: 13–30.
- Malarvizhi, A.; Kavitha, C.; Saravanan, M. and Ramesh, M. (2012). Carbamazepine (CBZ) induced enzymatic stress in gill, liver and muscle of a common carp, *Cyprinus carpio*. *J King Saud Univ*, 24: 179–186.
- McCord, J.M. and Fridovich, J. (1969). Superoxide dismutase: an enzymatic function for erythrocyte hemocuprein. *J Biol Chem*, 244: 6049–6055.
- Modesto, M.A. and Martinez, C.B.R. (2010). Roundup causes oxidative stress in liver and inhibits acetylcholine esterase in muscle and brain of the fish, *Prochilodus lineatus*. *Chemosphere*, 78: 294–299.
- Monteiro, D.A.; Rantin, F.T. and Kalinin, A.L. (2009). The effects of selenium on oxidative stress biomarkers in the freshwater characid fish matrinxã, *Brycon cephalus* exposed to organophosphate insecticide Folisuper 600 BR® (methyl parathion). *Comp Biochem Physiol C*, 149: 40–49.
- Muir, J. and Eduljee, G. (1999). PCP in the freshwater and marine environment of the European Union. *Sci Total Environ*, 236: 41–56.
- Nemcsok, J. and Boross, L. (1982). Comparative studies on the sensitivity of different fish species to metal pollution. *Acta Biol Hung*, 33: 23–27.
- Nwani, C.D.; Lakra, W.S.; Nagpure, N.S.; Kumar, R.; Kushwaha, B. and Srivastava, S.K. (2010). Toxicity of the herbicide atrazine: effects on lipid peroxidation and activities of antioxidant enzymes in the freshwater fish *Channa punctatus* (Bloch). *Int J Environ Res Public Health*, 7: 3298–3312.
- Office of Environmental Health Hazard Assessment (OEHHA) 1997. Inorganic Lead as a Toxic Air Contaminant. Part B. Health Effects of Airborne Inorganic Lead. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Air Toxicology and Epidemiology Section, Berkeley, CA.
- Osman, A.G.M.; Reheem, A.E.B.M.A.E.; AbuelFadl, K.Y. and GadEl-Rab, A.G. (2010). Enzymatic and histopathologic biomarkers as indicators of aquatic pollution in fishes. *Nat Sci*, 2: 1302–1311.
- Paglia, D.E. and Valentine, W.N. (1967). Studies on the Quantitative and Qualitative Characterization of Erythrocyte Glutathione Peroxidase. *Journal of Laboratory and Clinical Medicine*, 70: 158-169.
- Pampanin, D.M.; Camus, L.; Gomiero, A.; Marangon, I.; Volpato, E. and Nasci, C. (2005). Susceptibility to oxidative stress of mussels (*Mytilus galloprovincialis*) in the Venice Lagoon (Italy). *Mar Pollut Bull*, 50: 1548–1557.
- Pandey, A.K. and Dubey, S. (2015). Histological changes in liver and kidney of Catfish *H. fossilis* exposed to pentachlorophenol (PCP). *Plant Archives*, 15(2):1117-1120.
- Pandey, S.; Parvez, S.; Sayeed, I.; Haque, R.; Bin-Hafeez, B. and Raisuddin, S. (2003). Biomarkers of oxidative stress: a comparative study of river Yamuna fish *Walla goattu* (Bl. and Schn.). *Sci Total Environ*, 309: 105–115.
- Rao, V.J. (2006). Sublethal effects of an organophosphorus insecticide (RPR-II) on biochemical parameters of tilapia, *Oreochromis mossambicus*. *Comp Biochem Physiol C*, 143: 492–498.
- Ruas, C.B.; Carvalho Cdos, S.; de Araujo, H.S.; Espindola, E.L. and Fernandes, M.N. (2008). Oxidative stress biomarkers of exposure in the blood of cichlid species from a metal-contaminated river. *Ecotoxicol Environ Saf*, 71: 86–93.
- Sayeed, I.; Parvez, S.; Pandey, S.; Bin-Hafeez, B.; Haque, R. and Raisuddin, S. (2003). Oxidative stress biomarkers of exposure to deltamethrin in freshwater fish, *Channa punctatus* Bloch. *Ecotoxicol Environ Saf*, 56: 295–301.
- SEMA (Secretaria Estadual do Meio Ambiente) (1988). Avaliação da toxicidade de parapeixes (D.3). In Manual de Testes Para Avaliação da Ecotoxicidade de Agentes Químicos. Brasília, 10.
- Sen, I.; Shandil, A. and Shrivastava, V.S. (2011). Study for determination of heavy metals in fish species of the river Yamuna (Delhi) by inductively coupled plasma-optical emission spectroscopy (ICP-OES). *Adv. Appl. Sci. Res*, 2: 161–166.
- Sieja, K. and Talerczyk, M. (2004). Selenium as an element in the treatment of ovarian cancer in women receiving chemotherapy. *Gynecol Oncol*, 93: 320–327.
- Singh, M.; Sandhir, R. and Kiran, R. (2011). Effects on antioxidant status of liver following atrazine exposure and its attenuation by vitamin E. *Exp Toxicol Pathol*, 63: 269–276.
- Stoscheck, C.M. (1990). Quantitation of Protein. *Methods in Enzymology*, 182: 50-68.
- Sureda, A.; Batle, J.M.; Tauler, P.; Cases, N.; Aguiló, A.; Tur, J.A. and Pons, A. (2004). Neutrophil tolerance to oxidative stress induced by hypoxia/reoxygenation. *Free Radical Res*, 38: 1003–1009.
- Tripathi, G. and Shasmal, J. (2011). Concentration related responses of chlorpyrifos in antioxidant, anaerobic and protein synthesizing machinery of the freshwater fish, *Heteropneustes fossilis*. *Pestic Biochem Physiol*, 99: 215–220.
- Tyor, A. and Pahwa, K. (2017). Pollutants induced oxidative stress, DNA damage and cellular deformities in *Clarias gariepinus* (Burchell), from river Yamuna in Delhi region, India. *Bull Env Contam Toxicol*, 99: 33–38.
- Uner, N.; Oruc, E.O.; Sevgiler, Y.; Sahin, N.; Durmaz, H. and Usta, D. (2006). Effects of diazinon on acetylcholinesterase activity and lipid peroxidation in the brain of *Oreochromis niloticus*. *Environ Toxicol Pharmacol*, 21: 241–245.
- Vadhva, P. and Hasan, M. (1986). Organophosphate dichlorvos induced dose-related differential alterations in lipid levels and lipid peroxidation in various regions of the fish brain and spinal cord. *J Environ Sci Health B*, 21: 413–24.
- Van den Berg, K.J. (1990). Interaction of chlorinated phenols with thyroxine binding sites of Human transthyretin, albumin and thyroid binding globulin. *Chem Biol Interact*, 76: 63–75.

- Van der Oost, R., J. Beyer and Vermeulen, N.P.E. (2003). Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environ Toxicol Pharmacol*, 13: 57–149.
- Velkova-Jordanoska, L.; Kostoski, G. and Jordanoska, B. (2008). Antioxidative enzymes in fish as biochemical indicators of aquatic pollution. *Bulg J Agric Sci*, 14: 235-237.
- Verma, S.P. and Dubey, S. (2019). Histopathological changes in liver of *H. fossilis* exposed to pentachlorophenol (PCP). *Proceedings of Zoological Society of India*, 18(2): 103-108.
- Verma, S.P. and Dubey, S. (2021). Impact of Pentachlorophenol on histopathological changes in the testes of Catfish, *Heteropneustes fossilis*. *Proceedings of Zoological Society of India*, 20(1): 25-30.
- Vlahogianni, T.; Dassenakis, M.; Scoullou, M.J. and Valavanidis, A. (2007). Integrated use of biomarkers (superoxide dismutase, catalase and lipid peroxidation) in mussels *Mytilus galloprovincialis* for assessing heavy metals' pollution in coastal areas from the Saronikos Gulf of Greece. *Mar Pollut Bull*, 54: 1361–1371.
- Vutukuru, S.S.; Chintada, S.; Madhavi, K.R.; Rao, J.V. and Anjaneyulu, Y. (2006). Acute effects of copper on superoxide dismutase, catalase and lipid peroxidation in the freshwater teleost fish, *Esomus danricus*. *Fish Physiol Biochem*, 32: 221–229.
- Webb, D.; Gagnon, M.M. and Rose, T. (2005). Metabolic enzyme activities in black bream, *Acanthopagrus butcheri* from the swan canning estuary, Western Australia. *Comp Biochem Physiol C*, 141: 356–365.
- Wheatley, A.R. (2000). Some recent trends in the analytical chemistry of lipid peroxidation. *Trends Anal Chem*, 19: 617–628.
- WHO (1987). Pentachlorophenol. Geneva, World Health Organization, International Programme on Chemical Safety (Environmental Health Criteria 71).
- Winston, G.W. and Di Giulio, R.T. (1991). Prooxidant and antioxidant mechanisms in aquatic organisms. *Aquat Toxicol*, 19: 137–161.
- Zheng, G.J.; Leung, A.O.W.; Jiao, L.P. and Wong, M.H. (2008). Polychlorinated dibenzo-p-dioxins and dibenzofurans pollution in China: sources, environmental levels and potential human health impacts. *Environ Int*, 34: 1050–61.
- Zhou, X.N.; Wang, L.Y.; Chen, M.G.; Wu, X.H.; Jiang, Q.W.; Chen, X.Y.; Zhengand, J. and Utzinger, J. (2005). The public health significance and control of schistosomiasis in China-then and now. *Acta Trop*, 96: 97-105.