



Plant Archives

Journal homepage: <http://www.plantarchives.org>

DOI Url : <https://doi.org/10.51470/PLANTARCHIVES.2024.v24.no.2.156>

CORRELATION AND PATH ANALYSIS OF DIVERSE FENNEL (*FOENICULUM VULGARE* MILL.) GENOTYPES FOR SEED YIELD AND ASSOCIATED TRAITS

N. Divyashree^{1*}, Y.C. Vishwanath¹, S.C. Sarvamangala², B.N. Vijaykumar¹, H. Manjunath³, P. Bapurayagowda⁴ and K. Krishna⁵

¹Department of Plantation, Spices, Medicinal and Aromatic Crops, College of Horticulture, UHS, Bagalkot-587 104, Karnataka, India.

²Department of Genetics and Plant Breeding, College of Horticulture, UHS, Bagalkot - 587 104, Karnataka, India.

³Department of Plant Pathology, College of Horticulture, UHS, Bagalkot - 587 104, Karnataka, India.

⁴Department of Seed Science and Technology, College of Horticulture, UHS, Bagalkot - 587 104, Karnataka, India.

⁵Department of Agronomy, College of Horticulture, Munirabad, UHS, Bagalkot - 587 104, Karnataka, India.

*Corresponding author E-mail : divyashreenbalaji199095@gmail.com

(Date of Receiving-16-06-2024; Date of Acceptance-24-08-2024)

ABSTRACT

In the present investigation, thirty-eight diverse genotypes were utilized to understand the correlation and path analysis among seed yield and related yield attributing characteristics in fennel. The experiment was conducted following a randomized complete block design at the College of Horticulture, Bagalkot, during the year 2020-21. Data pertaining to 17 agronomic and quality traits were recorded in three replicates and subjected to statistical analysis. A majority of the traits exhibited high correlation coefficients at both genotypic and phenotypic levels, indicating inherent associations among them. Parameters such as plant height, days to initial flowering, days to maturity, number of umbels per plant, number of seeds per umbellet, and harvest index displayed significant positive correlations with seed yield per plant at both phenotypic and genotypic levels. Conversely, traits like dry umbel weight, test weight and aphid incidence showed negative associations with most of the studied traits. Based on path analysis, the number of branches per plant, days to 50 per cent flowering, number of seeds per umbellet, and oil content exerted a higher positive direct effect on seed yield per plant, revealing the need to emphasize on these characters for the genetic enhancement of fennel to augment seed yield.

Key words : Fennel, Seed yield, Correlation, Path analysis.

Introduction

Fennel (*Foeniculum vulgare* Mill), an essential seed spice, belongs to the Apiaceae family and is predominantly cultivated in temperate and subtropical regions for its seeds. With a rich history as one of the oldest seed spices, fennel offers a distinctive flavor utilized in various applications within the food, cosmetic and pharmaceutical industries due to its beneficial medicinal effects (Rather *et al.*, 2016). Research has highlighted the potential uses of the essential oil derived from fennel as a valuable antioxidant, antispasmodic, with antibacterial, anticancer and antifungal properties (El-Awadi and Hassan, 2010;

Badgajar *et al.*, 2014; Lucinewton *et al.*, 2005; Olle and Bender, 2010). The increasing commercial demand for fennel underscores the necessity to develop superior genotypes with enhanced yield and quality standards (Spice Board & Ministry of Agriculture and Farmers Welfare, Govt. of India, 2022-23).

Fennel originated from the Mediterranean region, where it exhibits a high level of genetic variability (Mirandli, 1999). Fennel breeding program plays a crucial role in its enhancement for cultivation and utilization. Commencing a breeding program, for any crop with genetic diversity necessitates acquiring information about agronomically

important traits to facilitate the selection and breeding of improved varieties effectively (Singh *et al.*, 2020).

The relationships between various traits and their effects on yield, both direct and indirect effects, lay the foundation for a successful breeding initiative (Kushwah *et al.*, 2021). As a quantitative trait, yield involves complex inheritance influenced by environmental fluctuations, thus demanding the indirect selection of highly heritable traits for yield enhancement (Terfa *et al.*, 2020). Additionally, understanding the associations and interactions of different traits with yield aids breeders in selection processes with enhanced precision and accuracy (Hassani *et al.*, 2023). The magnitude and direction of these trait associations with yield were assessed through genotypic and phenotypic correlation coefficients (Baye *et al.*, 2020). The determination of the relative importance of direct and indirect influences of the component characters towards seed yield is established through path analysis (Bhuva *et al.*, 2020). Breeding studies in various seed spices have utilized correlation and path analysis techniques (Kumar *et al.*, 2022). Therefore, detailed insights into significant characters for improvement programs can be obtained through correlation studies and path analysis. The primary objective of this study was to examine the direct and indirect influences of the component traits on seed yield and to identify key traits for consideration in fennel enhancement initiatives.

Materials and Methods

The research was conducted at Udyanagiri Campus, College of Horticulture, Bagalkot, Karnataka, India. Bagalkot situated in the northern dry zone of Karnataka, has co-ordinates of 16°12' N, 75°45' E, with an average elevation of around 610 m. The region experiences a warm and arid climate with minimal rainfall, averaging 318 mm annually, and is characterized by laterite soil.

From October 2020 to March 2021, the experiment took place to coincide with the cold and dry weather required by the fennel crop for flowering. A total of thirty-eight fennel genotypes were sourced from various institutions across India, showcasing a wide range of variability in 17 traits related to yield, including days to germination, plant height, number of branches per plant, days to first flowering, days to 50% flowering, days to maturity, number of umbels per plant, number of umbellets per umbel, number of seeds per umbellet, fresh umbel weight (g), dry umbel weight (g), test weight (g), biomass yield per plant (g), harvest index, oil content (%), aphid incidence and seed yield per plant (g). The experiment followed a Randomized Complete Block Design (RCBD) with two replications, with genotypes randomly assigned

within each replication.

The soil underwent two rounds of ploughing, harrowing, and was brought to a fine tilth prior to being divided into sections measuring 2.5 m × 2.2 m. In the process of land preparation, farmyard manure was administered as a basal dose at a rate of 15 tonnes per hectare, in conjunction with a complete dose of phosphorus and potassium, as well as a half dose of nitrogen (100:120:60 kg NPK/ha). At 30 days after sowing (DAS), the remaining half dose of nitrogen was applied as a top dressing. The crop was sown in lines in November 2020, with row spacing set at 50 cm. Thinning was carried out post emergence of seedlings by maintaining 20 cm between plants. Observations were recorded from five randomly selected plants in each treatment. Aphid incidence was documented during the flowering stage, as it was at its peak during this period. Upon reaching full maturity, the crop was harvested. Following the threshing process, the seeds underwent manual sorting and cleaning. Post field sampling from the seeds and subsequent drying of samples at 40°C, the essential oil was extracted through hydro-distillation in a Clevenger apparatus for approximately 3 hours. Each treatment involved the use of 100 g of dried seed powder and 1000 cc of water. The essential oils extracted from fennel seeds were obtained from dried green seeds (pasty stage) due to data indicating that fennel seeds at this stage possess the highest concentration of essential oil (Stefanini *et al.*, 2006).

Pearson's phenotypic and genotypic correlation coefficients were calculated for the 16 quantitative traits pairwise among the various cultivars as per the methodologies of Johanson *et al.* (1955), Miller *et al.* (1958), Singh and Chaudhury (1985). The statistical significance of the correlation coefficients was evaluated utilizing the table of "t" values by Fisher and Yates (1963) at (n-2) degrees of freedom. Among the seventeen phenotypic traits, seed yield was considered the dependent variable, while the remaining variables were viewed as independent. The standard genotypic path coefficients, equivalent to standard partial regression coefficients, were computed by solving a set of 'p' simultaneous equations using the "DOLITTLE TECHNIQUE" with the spar 1 computer software proposed by Goulden (1959).

The correlation coefficient may not always be sufficient to explain causal relationships between variables, as the association between two variables could be influenced by a third variable. Path analysis offers a rational interpretation of observed correlations by modelling the cause-and-effect relationships among

variables. Consequently, it becomes feasible to scrutinize the correlation coefficient of variables in terms of variance and covariance using path analysis (Gu *et al.*, 2019).

Path analysis has been developed as a statistical tool for analysing cause-and-effect relationships within a system of interconnected variables. A path coefficient, serving as a standardized partial regression coefficient, gauges the direct impact of a predictor variable on the dependent variable (Mohammadi *et al.*, 2003; Steel and Torrie, 1980). This facilitates the breakdown of the correlation coefficient into direct and indirect effects (Gu *et al.*, 2019), with unexplained effects being treated as residual effects. Plant breeders have utilized path analysis to comprehend the association between productivity and its constituents in different crops. In this study, the direct and indirect effects of path analysis were categorized using the Lenka and Misra (1973) scale.

Results and Discussion

Correlation study

A study of correlation was conducted to investigate the relationship between traits using Pearson's correlation coefficient. The correlation results are presented in Tables 1 and 2. Upon analysing genotypic and phenotypic correlations, it was found that there were positive and significant correlations of seed yield per plant with plant height ($r = 0.378^{**}$ and $r = 0.362^*$), days to first flowering ($r = 0.362^*$ and $r = 0.312^*$), number of umbels per plant ($r = 0.316^*$ and $r = 0.312^*$), number of seeds per umbellet ($r = 0.302^*$ and $r = 0.283^*$), dry umbel weight ($r = 0.325^*$ and $r = 0.319^*$), and harvest index ($r = 0.679^{**}$ and $r = 0.667^{**}$).

Table 1 : Pearson's genotypic correlation coefficient analysis among seed yield attributing quantitative traits of fennel genotypes.

Parameters	Genotypical Correlation Matrix																
	PH (cm)	NOB/P	DTG	DIFF	DT 50% F	DTM	NOU/P	NOU/U	NOF /Ut	FUW (g)	DUW (g)	TW (g)	BY/P (g)	HI	OC %	AI %	SY/P (g)
PH (cm)	1.0000	0.282*	0.341*	0.542**	0.0094	-0.360*	0.254*	0.434**	0.458**	0.407**	0.645**	-0.388**	0.0864	0.1134	0.1959	-0.0882	0.378**
NOB/P		1.0000	0.333*	0.384**	0.1132	-0.0880	0.248*	0.0186	-0.1605	0.671**	0.498**	-0.1894	0.0228	-0.0298	-0.308*	0.0008	-0.0782
DTG			1.0000	0.0625	0.331*	0.1987	0.1623	0.1720	0.1645	0.367*	0.378**	-0.256*	-0.0348	-0.0595	-0.1615	0.0488	-0.0697
DIFF				1.0000	0.412**	-0.368*	-0.0388	-0.0898	-0.0259	0.284*	0.454**	-0.288*	-0.1280	0.277*	-0.0914	-0.310*	0.362*
DT50% F					1.0000	0.410**	-0.280*	-0.1447	-0.1299	0.1754	0.0738	-0.1733	-0.1333	0.1584	-0.0211	0.0046	-0.0660
DTM						1.0000	-0.1357	-0.1755	-0.0355	-0.328*	-0.480**	0.1555	-0.0546	-0.1364	0.294*	0.1566	-0.367*
NOU/P							1.0000	0.278*	0.284*	0.347*	0.1979	-0.234*	0.2247	0.1415	-0.0839	-0.0580	0.316*
NOU/U								1.0000	0.713**	0.305*	0.390**	-0.2193	0.321*	-0.1239	0.323*	0.1553	0.2031
NOS/Ut									1.0000	0.1224	0.234*	-0.0509	0.1766	0.0715	0.1628	-0.0648	0.302*
FUW (g)										1.0000	0.766**	-0.386**	0.289*	-0.0972	-0.0594	0.0167	0.1466
DUW (g)											1.0000	-0.400**	0.1236	-0.0394	0.0079	0.325*	
TW (g)												1.0000	-0.321*	0.1445	0.0062	-0.2117	-0.290*
BY/P (g)													1.0000	-0.720**	0.395**	0.228*	-0.0935
HI														1.0000	-0.327*	-0.302*	0.679**
OC %															1.0000	0.0934	-0.1976
AI %																1.0000	-0.278*
SY/P (g)																	1.0000

*Significant @ 5% probability

** Significant @ 1% probability

PH: Plant height, NOB/P: Number of branches/ plant, DTG: Days to germination, DIFF: Days to first flowering, DT50%F: Days to 50% flowering, DTM: Days to maturity, NOU/P: Number of umbels/ plant, NOU/U: number of umbellets/ umbel, NOS/Ut: Number of seeds/ umbellet, FUW: Fresh umbel weight, DUW: Dry umbel weight, TW: Test weight, BY/P: Biomass yield/ plant, HI: Harvest index, AI: Oil content, OC: Oil content, AI: Aphid incidence, SY/P: Seed yield/ plant.

Table 2 : Pearson's phenotypic correlation coefficient analysis among seed yield attributing quantitative traits of fennel genotypes.

Phenotypical Correlation Matrix																	
Parameters	PH (cm)	NOB/P	DIG	DIFP	DT 50%F	DTM	NOU/P	NOU/U	NOF /Ut	FUW (g)	DUW (g)	TW (g)	BY/P (g)	HI	OC %	AI %	SY/P (g)
PH (cm)	1.0000	0.238*	0.315*	0.428**	-0.0175	-0.270*	0.249*	0.423**	0.412**	0.388**	0.616**	-0.366*	0.1125	0.1179	0.1880	-0.0726	0.362*
NOB/P		1.0000	0.273*	0.338*	0.0959	-0.1385	0.2140	0.0043	-0.1416	0.602**	0.458**	-0.1706	0.0243	-0.0399	-0.249*	-0.0136	-0.0584
DIG			1.0000	0.0561	0.282*	0.0925	0.1498	0.1556	0.1703	0.358*	0.363*	-0.239*	-0.0442	-0.0558	-0.1532	0.0495	-0.0633
DIFP				1.0000	0.291*	-0.297*	-0.0647	-0.1083	-0.0268	0.253*	0.389**	-0.272*	-0.1239	0.230*	-0.0745	-0.230*	0.312*
DT50%F					1.0000	0.1618	-0.228*	-0.1541	-0.1021	0.1275	0.0578	-0.0969	-0.1119	0.0897	-0.0220	-0.0180	-0.0393
DTM						1.0000	-0.0813	-0.0771	0.0096	-0.229*	-0.343*	0.1209	-0.0748	-0.0720	0.1720	0.1433	-0.296*
NOU/P							1.0000	0.272*	0.262*	0.340*	0.1880	-0.2247	0.2060	0.1416	-0.0809	-0.0582	0.312*
NOU/U								1.0000	0.672**	0.302*	0.380**	-0.2076	0.308*	-0.1117	0.299*	0.1520	0.1964
NOS/Ut									1.0000	0.1309	0.2207	-0.0118	0.1567	0.0641	0.1191	-0.0650	0.283*
FUW (g)										1.0000	0.755**	-0.365*	0.276*	-0.0940	-0.0702	0.0198	0.1426
DUW (g)											1.0000	-0.373**	0.1298	0.1179	-0.0376	0.0147	0.319*
TW (g)												1.0000	-0.281*	0.1215	-0.0206	-0.1978	-0.281*
BY/P (g)													1.0000	-0.676**	0.367*	0.2206	-0.0829
HI														1.0000	-0.303*	-0.278*	0.667**
OC %															1.0000	0.0762	-0.1729
AI %																1.0000	-0.267*
SY/P (g)																	1.0000

*Significant @ 5% probability

** Significant @ 1% probability

PH: Plant height, NOB/P: Number of branches/ plant, DIG: Days to germination, DIFP: Days to first flowering, DT50%F: Days to 50% flowering, DTM: Days to maturity, NOU/P: Number of umbels/ plant, NOU/U: number of umbellets/ umbel, NOS/Ut: Number of seeds/ umbellet, FUW: Fresh umbel weight, DUW: Dry umbel weight, TW: Test weight, BY/P: Biomass yield/ plant, HI: Harvest index, OC: Oil content, AI: Aphid incidence, SY/P: Seed yield/ plant.

Improvement in these traits could lead to an enhancement in seed yield, suggesting that positive selection should be applied to these traits for seed yield improvement. These findings align with previous studies by various researchers viz., Jindla *et al.* (1985), Bhandari and Gupta (1991), Agnihotri *et al.* (1997), Garg *et al.* (2003), Jain *et al.* (2003), Singh and Sastry (2005), Lal (2007), Cosge *et al.* (2009), Idhol *et al.* (2009), Dashora and Sastry (2011), Mangesha *et al.* (2013), Yadav *et al.* (2013), Sefidan *et al.* (2014), Singh *et al.* (2015b), Mohan *et al.* (2016) and Ram *et al.* (2017).

On the other hand, traits such as days to maturity ($r = -0.367^*$ and $r = -0.296^*$), test weight ($r = -0.386^{**}$ and $r = -0.365^*$) and aphid incidence ($r = -0.278^*$ and $r = -0.267^*$) exhibited significant negative correlations with seed yield per plant. Days to maturity can serve as a selection criterion for increasing seed yield if negative selection is practiced, indicating that a shorter period to 50% flowering results in higher seed yield per hectare. Similar observations were reported by previous researchers Jindla *et al.* (1985), Bhandari and Gupta (1991) and Yadav *et al.* (2013).

Path coefficient analyses

Path coefficient analysis was conducted in the current study to explore the relationships among fennel components. The analysis focused on genotypic associations and is presented in Table 3.

The path coefficient analysis at the genotypic level unveiled that the number of branches per plant ($r = 2.7229$) had the highest and significant positive direct effect on seed yield per plant, followed by days to 50% flowering ($r = 2.3168$),

Table 3 : Genotypic path coefficient analysis among seed yield attributing quantitative traits of fennel genotypes.

Parameters	Genotypic path matrix of Seed yield/plant																
	PH (cm)	NOB/P	DIG	DIFF	DT 50%F	DTM	NOU/P	NOU/U	NOF /Ut	FUW (g)	DUW (g)	TW (g)	BY/P (g)	HI	OC %	AI %	SY/P (g)
PH (cm)	-0.8931	-0.2520	-0.3044	-0.4843	-0.0084	0.3211	-0.2271	-0.3871	-0.4089	-0.3632	-0.5759	0.3460	-0.0771	-0.1013	-0.1750	0.0788	0.378**
NOB/P	0.7685	2.7229	0.9057	1.0460	0.3082	-0.2398	0.6752	0.0505	-0.4369	1.8265	1.3565	-0.5158	0.0622	-0.0811	-0.8397	0.0022	-0.0782
DIG	0.0630	0.0614	0.1847	0.0115	0.0611	0.0367	0.0300	0.0318	0.0304	0.0678	0.0699	-0.0472	-0.0064	-0.0110	-0.0298	0.0090	-0.0697
DIFF	-1.2555	-0.8894	-0.1446	-2.3153	-0.9528	0.8509	0.0899	0.2080	0.0600	-0.6572	-1.0502	0.6662	0.2963	-0.6417	0.2115	0.7187	0.362*
DT50%F	0.0217	0.2622	0.7670	0.9535	2.3168	0.9502	-0.6476	-0.3353	-0.3009	0.4063	0.1709	-0.4015	-0.3089	0.3669	-0.0489	0.0108	-0.0660
DTM	1.2350	0.3025	-0.6827	1.2625	-1.4090	-3.4354	0.4663	0.6028	0.1219	1.1258	1.6482	-0.5343	0.1877	0.4687	-1.0092	-0.5380	-0.367*
NOU/P	0.1357	0.1323	0.0866	-0.0207	-0.1491	-0.0724	0.5336	0.1481	0.1515	0.1852	0.1056	-0.1249	0.1199	0.0755	-0.0448	-0.0309	0.316*
NOU/U	-0.7374	-0.0316	-0.2926	0.1528	0.2462	0.2985	-0.4722	-1.7012	-1.2136	-0.5195	-0.6627	0.3730	-0.5454	0.2108	-0.5499	-0.2642	0.2031
NOS/Ut	0.9596	-0.3363	0.3448	-0.0543	-0.2722	-0.0744	0.5950	1.4953	2.0960	0.2565	0.4896	-0.1067	0.3703	0.1499	0.3412	-0.1359	0.302*
FUW (g)	-1.1068	-1.8256	-0.9995	-0.7726	-0.4773	0.8919	-0.9448	-0.8311	-0.3331	-2.7215	-2.0849	1.0506	-0.7864	0.2645	0.1616	-0.0455	0.1466
DUW (g)	0.3180	0.2457	0.1865	0.2237	0.0364	-0.2366	0.0976	0.1921	0.1152	0.3778	0.4931	-0.1974	0.0610	0.0611	-0.0194	0.0039	0.325*
TW (g)	0.3875	0.1895	0.2557	0.2878	0.1733	-0.1556	0.2340	0.2193	0.0509	0.3861	0.4004	-1.0002	0.3214	-0.1445	-0.0062	0.2117	-0.290*
BY/P (g)	-0.0129	-0.0034	0.0052	0.0191	0.0199	0.0082	-0.0336	-0.0479	-0.0264	-0.0431	-0.0185	0.0480	-0.1493	0.1075	-0.0590	-0.0340	-0.0935
HI	0.0617	-0.0162	-0.0323	0.1507	0.0861	-0.0742	0.0769	-0.0674	0.0389	-0.0528	0.0673	0.0786	-0.3914	0.5436	-0.1779	-0.1642	0.679**
OC %	0.4066	-0.6400	-0.3352	-0.1896	-0.0438	0.6097	-0.1741	0.6709	0.3379	-0.1233	-0.0818	0.0129	0.8197	-0.6793	2.0754	0.1938	-0.1976
AI %	0.0260	-0.0002	-0.0144	0.0914	-0.0014	-0.0461	0.0171	-0.0457	0.0191	-0.0049	-0.0023	0.0623	-0.0671	0.0889	-0.0275	-0.2945	-0.278*
SY/P (g)	0.378**	-0.0782	-0.0697	0.362*	-0.0660	-0.367*	0.316*	0.2031	0.302*	0.1466	0.325*	-0.290*	-0.0935	0.679**	-0.1976	-0.278*	1.0000
Partial R ²	-0.3373	-0.2130	-0.0129	-0.8386	-0.1530	1.2617	0.1688	-0.3456	0.6330	-0.3990	0.1604	0.2903	0.0140	0.3688	-0.4102	0.0820	

*Significant @ 5% probability

** Significant @ 1% probability

Residual effect: 0.855

PH: Plant height, NOB/P: Number of branches/ plant, DTG: Days to germination, DTFF: Days to first flowering, DT50%F: Days to 50 % flowering, DTM: Days to maturity, NOU/P: Number of umbels/ plant, NOU/U: number of umbellets/ umbel, NOS/Ut: Number of umbellets/ umbellet, FUW: Fresh umbel weight, DUW: Dry umbel weight, TW: Test weight, BY/P: Biomass yield/ plant, HI: Harvest index, OC: Oil content, AI: Aphid incidence, SY/P: Seed yield/ plant




Fig. 1 : Genotypical path diagram for seed yield per plant from their associated traits among 38 fennel genotypes.

number of seeds per umbellet ($r = 2.0960$), and oil content ($r = 2.0754$). Conversely, harvest index ($r = 0.5436$) and number of umbels per plant ($r = 0.5336$) showed non-significant positive direct effects on seed yield per plant. Although these traits had significant correlations with seed yield per plant, indicating indirect cumulative effects on seed yield through other characters. High indirect effects of these characteristics were predominantly observed via days required for maturity, seed weight, and biomass yield per individual plant, underscoring the significance of these traits as pivotal selection criteria for enhancing seed yield in fennel. These findings align with prior studies by Bhandari and Gupta (1991), Agnihotri *et al.* (1997), Jain *et al.* (2003), Dashora and Sastry (2011), Yadav *et al.* (2013), and Sefidan *et al.* (2014).

Meanwhile, days to maturity ($r = -3.4354$), weight of fresh umbels ($r = -2.7215$), days to initial flowering ($r = -2.3153$), number of umbellets per umbel ($r = -1.7012$), test weight ($r = -1.0002$), plant height ($r = -0.8931$) and aphid infestation ($r = -0.2945$), exhibited sequentially declining negative direct impacts on seed yield per plant. Similar outcomes were documented by Agnihotri *et al.* (1997), Jain *et al.* (2003), Mangesha *et al.* (2013) and Yadav *et al.* (2013).

The residual effects depicted the extent to which the causal factors elucidated the variability in the dependent factor. Minimal residual effects signified that the causal factors adequately explicated the dependent variable. In this investigation, the residual effect for seed yield was computed to be 0.855. As the residual effects for seed

yield were relatively substantial, it is plausible that there exist essential additional factors influencing seed yield that were not accounted for in this study. The path diagram illustrating seed yield is presented in Fig. 1.

It is deduced that a substantial level of association exists among 17 traits of fennel (*Foeniculum vulgare* Mill.), as evidenced by the findings of the correlation and path coefficient analysis. The genotypic and phenotypic assessment demonstrated a significant and positive correlation between seed yield and various traits such as plant height, days to initial flowering, days to maturity, number of umbels per plant, number of seeds per umbellet and harvest index. The results of the path analysis highlighted those key variables including the count of seeds per umbellet, number of branches per plant, days to reach 50 per cent flowering and oil content were the major contributors to seed yield. Additionally, there was an indication that the number of umbels per plant and harvest index also played a role in influencing seed yield.

Acknowledgment

I sincerely acknowledged University of Horticultural Sciences, Bagalkot for willingly provided the land for the research work at the campus, and my major advisor (Dr. Y.C. Vishwanath) and committee members (Dr. Sarvamangala. S. Cholin, Dr. Vijaykumar. B. Narayanpur, Dr. Manjunath Hubballi, Dr. Baburayagowda Patil and Dr. Krishna Kurubetta) for their financial, resources and knowledge assistance towards my project. Dr. Sarvamangala S. Cholin (Co-guide) is acknowledged for provided facilities for laboratory analysis and assisted with statistical analysis. Dr. Krishna Kurubetta for providing genetic resources for my project. I also thank my field assistant, field workers, my friends and juniors for their assonance in field work and laboratory analysis.

References

- Agnihotri, P., Dashora S.L. and Sharma R.K. (1997). Variability, correlation, and path analysis in fennel (*Foeniculum vulgare* Mill.). *J. Spices Arom. Crops*, **6**(1), 51-54.
- Badgujar, S.B., Patel V.V. and Bandivdekar A.H. (2014). *Foeniculum vulgare* Mill.: A review of its botany, phytochemistry, pharmacology, contemporary application and toxicology. *Biomed Res. Int.*, 1-32.
- Bagheri, N., Babaeian-Jelodar N. and Pasha A. (2011). Path coefficient analysis for yield and yield components in diverse rice (*Oryza sativa* L.) genotypes. *Biharean Biologist*, **5**(1), 32-35.
- Bahmani, K., Izadi-Darbandi A., Noori S.A.S., Jafari A.A. and Moradi N. (2012). Determination of interrelationships among phenotypic traits of Iranian fennel (*Foeniculum vulgare* Mill.) using correlation, stepwise regression, and path analyses. *J. Essen. Oil Bear. Plants*, **15**(3), 424 - 444.
- Baye, A., Berihun B., Bantayehu M. and Derebe B. (2020).

- Genotypic and phenotypic correlation and path coefficient analysis for yield and yield-related traits in advanced bread wheat (*Triticum aestivum* L.) lines. *Cog. Food Agri.*, **6(1)**, 1-17.
- Bhandari, M.M. and Gupta A. (1991). Variation and association analysis in coriander. *Euphy.*, **58(1)**, 1-4.
- Bhuva, R.B., Babariya C.A., Movaliya H.M., Gadhiya J.A. and Balar V.S. (2020). Correlation and path analysis for seed yield in soybean [*Glycine max* (L.) Merrill.]. *Indian J. Pur. App. Bios.*, **8(4)**, 375-380.
- Dashora, A. and Sastry E.V.D. (2011). Variability, character association and path coefficient analysis in fennel. *Indian J. Horti.*, **68(3)**, 351-356.
- Dewey, D.R. and Lu K.N. (1959). A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron. J.*, **51**, 515-518.
- El-Awadi, M.E. and Hassan E.A. (2010). Physiological responses of fennel (*Foeniculum vulgare* Mill.) plants to some growth substances: The effect of certain amino acids and a pyrimidine derivative. *J. Am. Sci.*, **6(7)**, 120-125.
- Fisher, R.A. and Yates F. (1963). Statistical tables for biological agricultural and medical research. *Oliv. Edinb.*, 146.
- Goulden, C. (1959). *Methods of statistical analysis*. Asia Publishing House, Calcutta, India.
- Gu, F., Yung Y.F. and Cheung M.W.L. (2019). Four covariance structure models for canonical correlation analysis: a COSAN modelling approach. *Multiv. Beh. Res.*, **54(2)**, 192-223.
- Hassani, M., Mahmoudi S.B., Saremrad A. and Taleghani D. (2023). Genotype by environment and genotype by yield and trait interactions in sugar beet: analyzing yield stability and determining key traits association. *Scientific Reports*, **13(1)**, 1-16.
- Jain, U.K., Singh D. and Amrita (2003). Correlation and path analysis for certain metric traits in coriander. *Progre. Agri.*, **3(1&2)**, 86-88.
- Jindla, L.N., Singh T.N., Rangalla H. and Bansal M.L. (1985). Genetic variability and path coefficient analysis in coriander. *Crop Impro.*, **12(2)**, 133-136.
- Johanson, H.W., Robinson H.F. and Comstock R.E. (1955). Estimates of genetic and environmental variability of soybean. *Agron. J.*, **47**, 314-318.
- Kumar, M.A., Choudhary G., Garhwal O.P. and Netwal M. (2022). Correlation coefficient and path analysis for yield traits in coriander (*Coriandrum sativum* L.) genotypes. *Elect. J. Plant Breed.*, **13(1)**, 253-257.
- Kushwah, A., Sikarwar R.S., Vishwakarma M., Tiwari S. and Singh S. (2021). Relationship among the yield and its yield contributing traits in spring wheat (*Triticum aestivum* L.). *Phar. Inno. J.*, **10(9)**, 1974-1977.
- Lenka, D. and Misra B. (1973). Path coefficient analysis of yield in rice varieties. *Indian J. Agric. Sci.*, **43**, 376-379.
- Lucinewton, S., Raul N., Carvalho J., Mirian B., Lin C. and Angela A. (2005). Supercritical fluid extraction from fennel (*Foeniculum vulgare* Mill.) global yield, composition, and kinetic data. *J. Supercri. Flu.*, **35**, 212-219.
- Mangesha, B., Alemaw G and Tesfaye B. (2013). Correlation studies and path coefficient analysis for seed yield and yield components in Ethiopian coriander accessions. *Afr. Crop Sci. J.*, **21(1)**, 51-59.
- Miller, P.A., Williams J.C. and Comstock H.F. (1958). Estimation of genetic and environmental variance in upland cotton and their implications in selection. *Agron. J.*, **50**, 126-131.
- Miranldi, E. (1999). Comparison of the essential oil from ten *Foeniculum vulgare* Mill. samples of fruits of different origin. *Flav. Frag. J.*, **14**, 379-382.
- Mohammadi, S.A., Prasanna B.M. and Singh N.N. (2003). Sequential path model for determining interrelationship among grain yield and related characters in maize. *Crop Sci.*, **43**, 1690-1697.
- Mohan C., Singh J.P., Kumar S. and Puri A. (2016). Studies on character association in fennel (*Foeniculum vulgare* Mill.). *Int. J. Plant Soil Sci.*, **14(6)**, 1-5.
- Olle, M. and Bender I (2010) The content of oils in umbelliferous crops and its formation. *Agron. Res.*, **8 (Special Issue III)**, 687-696.
- Ram, H., Khan M.M., Pandey V.P. and Dwivedi D.K. (2017). Study on correlation coefficient and path analysis in coriander (*Coriandrum sativum* L.) genotypes. *Indian J. Biotechn.*, **9(2)**, 332-338.
- Rather, M.A., Dar B.A., Sofi S.N., Bhat B.A. and Qurishi M.A. (2016). *Foeniculum vulgare*: A comprehensive review of its traditional use, phytochemistry, pharmacology and safety. *Arabian J. Chem.*, **9**, 1574-1583.
- Sefidan, A. Y., Valizadeh M., Aharizad S. and Sabzi M. (2014). Path analysis of grain yield, some morphological traits and essential oil content in different fennel (*Foeniculum vulgare* Mill.) populations. *J. Biodi. Envi. Sci.*, **4(5)**, 10-15.
- Singh, B., Singh K.P. and Sengupta S.K. (2015b). Correlation analysis traits in elite genotypes of coriander. *J. Breed. Genet.*, **47(1)**, 70-78.
- Singh, R.K. and Chandhury B.D. (1985). *Biometrical methods in quantitative genetic analysis*. Kalyni Publishers, New Delhi, India. pp. 318.
- Singh, R.K., Prasad A., Muthamilarasan M., Parida S.K. and Prasad M. (2020). Breeding and biotechnological interventions for trait improvement: status and prospects. *Planta* **252**, 1-18.
- Spice Board & Ministry of Agriculture and Farmers Welfare, *Govt. of India* (2022-23).
- Steel, R.G.D. and Torrie J.H. (1981). *Principles and procedures of statistics, a biometrical approach*. McGraw Hill, New York, NY.
- Stefanini, M.B., Ming L.C., Marques M.O.M., Facanali R., Meireles M.A.A., Moura L.S., Marchese J.A. and Sousa L.A. (2006). Essential oil constituents of different organs of fennel (*Foeniculum vulgare* var. *vulgare*). *Rev. Bras. Pl. Med., Botucatu.*, **8**, 193-198.
- Terfa, G.N. and Gurmu G.N. (2020). Genetic variability, heritability and genetic advance in linseed (*Linum usitatissimum* L) genotypes for seed yield and other agronomic traits. *Oil Crop Sci.*, **5(3)**, 156-160.
- Wright, S. (1921). Correlation and causation. *J. Agric. Res.*, **20**, 557-585.
- Yadav, P.S., Pandey V.P. and Yadav Y. (2013). Variability studies in fennel (*Foeniculum vulgare* Mill.). *J. Spic. Arom.*, **22(2)**, 5-11.