



# Plant Archives

Journal homepage: <http://www.plantarchives.org>  
DOI Url : <https://doi.org/10.51470/PLANTARCHIVES.2022.v22.no1.006>

## ESSENTIAL OIL FROM AERIAL PARTS OF *RHETINOLEPIS LONADIOIDES* (COSS.): EXTRACTION, CHEMICAL COMPOSITION AND ANTIMICROBIAL ACTIVITY

Missoum Mansour<sup>1,2</sup>, Gherib Mohammed<sup>3\*</sup>, Ziane Mohammed<sup>1,4\*</sup>, Ben Braïek Olfa<sup>5</sup>,  
Yousfi Mohammed<sup>6</sup> and Amrouche Abdel Ilah<sup>3</sup>

<sup>1</sup>Université de Ain Témouchent, route de Sidi bel abbès, N101, Ain Témouchent 46000, Algérie.

<sup>2</sup>Laboratoire d'hydrologie appliquée et environnement, Université de Ain Témouchent, Algérie.

<sup>3</sup>Laboratoire Gestion durable des ressources naturelles dans des zones arides et semi-arides, Département de SNV, Institut des sciences et technologies, Centre Universitaire Salhi Ahmed 45000 Nâama Algérie

<sup>4</sup>Laboratoire de Microbiologie Appliquée à l'Agroalimentaire, au Biomédical et à l'Environnement (LAMAABE), Faculté des SNV/STU, Université de Tlemcen 13000 Tlemcen, Algérie

<sup>5</sup>Laboratory of Transmissible Diseases and Biologically Active Substances (LR99ES27), Faculty of Pharmacy, University of Monastir, Tunisia

<sup>6</sup>Université Amar Telidji Laghouat Département des Sciences de la matière

(Date of Receiving : 15-08-2021; Date of Acceptance : 06-11-2021)

### ABSTRACT

In this study, the chemical composition and antimicrobial activity of the essential oil (EO) from aerial parts of *Rhetinolepis lonadioides* collected from southwestern Algeria were assessed. The EO was obtained by hydrodistillation and analyzed by GC/MS. In total, thirty-two compounds were identified representing 80.00% of the total of the EO, with  $\beta$ -Pinene (41.85%) being the major monoterpenes. Phenanthroquinone (10.39%) and Ethanone, 1-[1,1'-biphenyl]-4-yl- (2.66%) were the main oxygenated monoterpenes, and the diterpene hydrocarbons were mainly represented by m-Camphorene (5.43%). Six EO samples of *R. lonadioides* collected in location exhibited similar chemical composition evidencing a chemical homogeneity. The agar disc diffusion method showed that *R. lonadioides* EO was effective against *Bacillus cereus*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* with diameters of inhibition ranging from  $17 \pm 2.65$  to  $12.33 \pm 1.13$  mm, and the minimum inhibitory concentration (MIC) values were between 4 and 10  $\mu$ L/mL. The lowest MIC values (4.0  $\mu$ L/mL) were measured for *Bacillus cereus*.

**Keywords:** antimicrobial activity, chemical composition, essential oil, GC/MS, *Rhetinolepis lonadioides*;

### Introduction

Essential oils (EOs) and volatile constituents extracted from aromatic plants are frequently used for diverse purposes especially in folk medicine for prevention and treatment of different human diseases, such as bacterial and viral infections (Rates 2001). Indeed, they have gained an increasing interest in these last years in pharmaceutical and food industries due to their safety aspects as they are natural products. Hence, many investigations of new active EOs for such applications are intensively conducted.

The species *Rhetinolepis lonadioides* (*R. lonadioides*), also known as *Ormenis lonadioides* (Coss.) Maire, or *Anthemis lonadioides* (Coss.) Hochr. is endemic to Algeria and Morocco (Oberprieler, 1998). *Anthemis* L. (*Asteraceae*) is one of the largest genus of the tribe *Anthemideae* which consists of nearly 175 species worldwide. It is distributed widely across Europe, South-West Asia, North and East Africa (Oberprieler *et al.*, 2007). South-West Asia is one of the genetic centers of the genus *Anthemis* (Rates, 2001). In fact, *R. lonadioides* is an annual plant, finely canescent pubescent, of 10-20 cm high, branched from the base with stiff stems, erect or lying with ascending hairy leaves with a

blade progressively thinned into a petiole and 3-5 lobed at the apex, with short subtute segments, and the bracts are entire and superior. It is characterized by a synflorescence in loose corymbs, discoid heads, obconic (diameter 5-8 mm), homogamous, on a short peduncle (3-12 mm), involucre with bracts in 1-2 rows, convex receptacle with narrow paleas, acuminate and hairy. Its flowers are tubular ( $\leq 3$  mm) and yellow, with enlarged subailed tube and 5-lobed corolla. Its achenes are less than 2mm, triangular, compressed, glabrous, bald, with numerous fine longitudinal striations and a central vein more or less marked on the ventral surface (Battandier 1888, African Plant Database, 2021).

*Anthemis* species have several biological activities and are widely used in folk medicine for treatment of gastrointestinal disorders, haemorrhoids, cough, stomach aches and liver failure (Baytop, 1999; Kultur, 2007; Ugurlu and Secmen 2008; Gonenc *et al.*, 2011; Korkmaz and Karakus, 2015). In addition, they are able to soothe pains and irritations, clean wounds (Pavlovic *et al.*, 2010), used as herb teas, and used in cosmetics as well as in the pharmaceutical industry (Kivcak *et al.*, 2007). Several *Anthemis* spp. have been studied for their essential oils (Uzel *et al.*, 2004;

Javidnia *et al.*, 2004; Kurtulmus *et al.* 2009; Yusufoglu *et al.*, 2018), secondary metabolites, terpenoids, sesquiterpene lactones, flavonoids and coumarins (Hofer and Greger 1985; Bruno *et al.*, 1997; Vajs *et al.*, 1999; Gonenc *et al.*, 2011; Tawaha *et al.*, 2015; Orlando *et al.*, 2019) and their neuro protective (Venditti *et al.*, 2016), cytotoxic (Alessandro *et al.*, 2016), antioxidant and antimicrobial effects (Kivcak *et al.*, 2007; Albayrak and Aksoy, 2012; Stojkovic *et al.*, 2014; Guragaç Dereli *et al.*, 2018).

To our knowledge, there are about twenty publications in the literature reporting essential oils from the aerial parts of 17 species and subspecies of plants of the *Anthemis* genus. None concerns *R. lonadioides* (Coss.).

Also, no research has yet been carried out to determine the chemical composition of the essential oil from aerial parts of *R. Lonadioides* or to evaluate its antimicrobial activity. For this reason, our study was undertaken to assess, for the first time, the chemical composition of Algerian essential oil extracted from *R. lonadioides*. Then, the characterized *R. lonadioides* essential oil was evaluated for its *in vitro* antimicrobial effect against some microbial pathogens. The expected results were then used for the valorization of *R. lonadioides* essential oil as a new bio agent for several pharmaceutical or food applications.

## Material and Methods

### Plant Material

Samples of aerial parts of *R. lonadioides* were collected during the flowering period (April, 2021) in the region of Beni Abbes (southwest of Algeria 30° 05' north, 2° 06' west). A voucher specimen has been deposited at the Laboratory Sustainable management of natural resources in arid and semi-arid zones, Department of SNV, Institute of Science and Technology, University Center Salhi Ahmed Nâama Algeria. Dried aerial parts (around 300–400 g) were submitted to hydrodistillation, using a Clevenger-type apparatus for 3 hours. Yields have been calculated from dry material.

### Gas Chromatography (GC) Analysis

GC analyses were performed with a BrukerScion SQ. It is equipped with a DB-5 type capillary column (length: 25 m, internal diameter: 0.220 mm, film thickness: 0.25 µm). The column temperature was set at 50°C for 10 min, then raised gradually from 50 to 250°C at a rise rate of 2°C/min, and finally it was maintained at 250°C for 15 min. The injector temperature was set at 250°C. The injection mode was Split (division ratio of 1:100). The carrier gas flow rate (helium) was set at 1 mL/min. the volume of the injected sample was 0.2 µL.

### Gas chromatography-mass spectroscopy (GC/MS) analysis

The analysis of the *R. lonadioides* essential oil samples was carried out in the Technical Platform of Physico-Chemical Analysis (PTAPC-CRAPC)-Laghout-Algeria, using a SHIMADZU GCMSQP2020 Instruments, equipped with a fused Rxi®-5ms capillary column (30 m × 0.25 mm, 0.25 µm film thickness, Phase: Cross bond® 5% diphenyl /95% dimethylpolysiloxane). A solution of 0.5 µL solution prepared by 10% vol. of the sample dilution in n-hexane, was injected in split mode (30:1). Injector and detector temperatures were maintained at 250°C and 310°C, respectively. The column temperature was programmed at 60°C for 3 min then increased gradually to 310°C at 2°C/min,

and finally maintained at 310°C for 10 min. The carrier gas used was Helium (99.995% purity) with a flow rate of 1 mL/min. The mass spectrometer conditions were as follows: ionization voltage 70eV, ion source temperature 200°C, and electron ionization mass spectra were acquired over the mass range of 45–600 m/z.

### Identification of Essential Oil Composition

Identification of components was performed by comparing their relative retention index (RI) determined with the reference of homologous series of *n*-alkanes (C8 to C24) (Rustaiyan *et al.*, 2004 and Toiu *et al.*, 2009).

The fragmentation patterns of the mass spectra were compared with the WILEY and NIST 05 libraries. The linear temperature-programmed RIs of all the constituents were calculated based on the GC through the interpolation between bracketing *n*-alkanes as follows:

$$RI = 100 \times \left[ \frac{t_{R(i)} + t_{R(z)}}{t_{R(z+1)} + t_{R(z)}} Z \right]$$

Where Z: the number of carbon atoms in the smaller *n*-alkane.

$t_{R(i)}$ ,  $t_{R(z)}$  and  $t_{R(z+1)}$ : the retention time of the desired compound, the smaller *n*-alkane and the larger *n*-alkane, respectively (Jalali-Heravi *et al.*, 2015).

### Antimicrobial Activity of the Essential Oil

#### Microbial Strains

Antimicrobial activity of the essential oil extracted from the aerial parts of *R. lonadioides* was evaluated against five Gram-positive bacteria (*Bacillus cereus* ATCC 10876, *Bacillus subtilis* ATCC 21332, *Listeria monocytogenes* ATCC 15313, *Staphylococcus aureus* ATCC 25923, and *Enterococcus faecalis* ATCC 49452) and two Gram-negative bacteria (*Escherichia coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC27853).

#### Screening for Antimicrobial Activity

The agar diffusion method (NCCLS 1997) was used for the determination of the antimicrobial activity of the essential oils. Briefly, 1 mL of the tested microbial suspension (at 10<sup>6</sup> cells/mL) was spread onto the surface of Mueller Hinton Agar (Biokar diagnostics, Beauvais, France) plates. The *R. lonadioides* essential oil was dissolved in dimethyl sulfoxide (DMSO) at 10% to enhance oil solubility. Then, filter paper discs (6 mm of diameter) were impregnated with the prepared mixture of DMSO and *R. lonadioides* essential oil and placed on the surface of the inoculated plates that were afterwards incubated at 37°C for 24 hours. After incubation, observed clear zones of inhibition were measured (mm). DMSO and antibiotics [spiramycin 100 µg and Oflatoxin 5 µg, BioMérieux] were used as controls. Each test was performed in triplicate in at least three separate experiments.

#### Determination of Minimum Inhibitory Concentration (MIC)

The MIC values of the *R. lonadioides* essential oil were determined by the micro dilution assay in 96-well microplates. The essential oil was dissolved in DMSO then it was first diluted to the highest concentration, followed by serial dilutions made in Mueller-Hinton broth (Biokar diagnostics, Beauvais, France) over the concentration range 1–6 µL/mL. Ten µL of standardized suspension was added.


Inoculated plates were incubated at 37°C for 24 hours for the bacteria. The MIC was defined as the lowest concentration of *R. lonadioides* essential oil resulting in complete inhibition of visible growth after incubation (Ahmed Elkhalfifa *et al.*, 2018). Tests were performed three times in triplicate.

### Statistical Analysis

All the experimental results were submitted for a variance analysis (ANOVA) using the SPSS software (version 11.5). Means and standard errors were calculated and a probability level of  $P < 0.05$  was used in testing the statistical significance of all the data. Tukey's post hoc test was used to determine significance of mean values for multiple comparison at  $P < 0.05$ .

## Results and Discussion

Essential oil samples were obtained by the hydrodistillation of dry aerial parts of *R. lonadioides* Coss. Yields of essential oil, calculated w/w versus dry material, ranged between 0.90% and 1.40%. These obtained EO yields are greater than those found for some plants of the same *Anthemis* genus such as *Anthemis nobilis* L. with an EO yield value of 0.67% (Sadiki and Idrissi 2019), *Anthemis maritima* from Corsica (0.015%) (Collu *et al.*, 2008) and *Anthemis cretica* sub sp. *carpatica* (0.22%) (Kürkçüoğlu and Tosun, 2020). Indeed, the yield of an EO depends on many factors (growth stage, pedoclimatic conditions, extraction technique, etc.). One oil sample was analyzed by GC(RI) and GC/MS (Table 1, Fig. 1).



**Fig. 1 :** Gas chromatogram GC/SM of *Rhadinolepis lonadioides* Coss. essential oil The numbered peaks are the identified components as listed in Table 1

### Chemical Composition of *R. lonadioides* (Coss.) Essential Oil

The detailed analysis of a bulk sample was carried out by a combination of chromatographic and spectroscopic techniques. One oil sample was analyzed by GC(RI) and GC(MS) (Table 1). In total, thirty-two compounds were identified, accounting for 80.00% of the whole composition, and were dominated by monoterpenes (68.33%). The hydrocarbon monoterpenes (53.94%) fraction was mostly represented by  $\beta$ -Pinene (41.85%), D-Limonene (6.79%),  $\alpha$ -Fenchene (3.64%) and  $\alpha$ -Pinene (1.01%). Hydrocarbons sesquiterpene (7.99%) were well represented by (E)-beta-Famesene (3.19%) and  $\alpha$ -Curcumene (1.63%), while, m-Camphorene (5.43%) was the only diterpene hydrocarbons present in a significant ( $P < 0.05$ ) amount. Moreover, this essential oil is characterized by the presence of oxygenated sesquiterpene of, Eudesm-7(11)-en-4-ol (0.47%) and Nerolidol (0.17%).

According to our results, the composition of the essential oil isolated from Algerian *R. lonadioides* differs drastically ( $P < 0.05$ ) from those of other *Anthemis* species dominated by oxygenated compounds, *A. montana* ( $\alpha$ -Thujone 46.9%,  $\beta$ -Thujone 16%, Trans chrysanthemyl

11.3%) (Bulatovic *et al.*, 1998), *A. tinctoria* L. (1,8 cineole 7.9%,  $\beta$ -Pinene 7.3%,  $\alpha$ -Pinene 4.4%) (Holla *et al.*, 2002) and *A. talyshensis* A. ( $\alpha$ -Eudesmol 18.2%, Borneol 13.3%, Hexadecanoic acid 9.5%) (Aghajani *et al.*, 2005). On the other hand, it presents some analogies with *Anthemis* essential oils that contained appreciable contents of monoterpene hydrocarbons: 2,4-Thujadiene,  $\beta$ -Pinene or Paracymene. For instance, *A. melampodina* (Grace, 2002) contained  $\beta$ -Pinene (6.4%) and Paracymene (11%), *A. altissima* (Javidnia *et al.*, 2004) (2,4-Thujadiene at 27% and  $\alpha$ -Pinene at 4%) and *A. melanolepis* (Saroglou *et al.*, 2005) ( $\beta$ -Pinene 11.7%).

Generally, the difference in chemical composition of our essential oil comparing to others from different *Anthemis* plants could be attributed to many factors such as the plant species, the part of the plant from which was extracted the essential oil, the genetic factors as well as the physiological, environmental and geographic variations (Dridi *et al.*, 2020).

### Antimicrobial Activity

The antimicrobial activity of *R. lonadioides* EO was assayed against seven bacteria using the agar disc diffusion method and measuring the minimum inhibitory concentration (MIC) (Table 2). The results showed that the EO had

substantial antimicrobial activity. In fact, it was effective against *B. cereus* and *P. aeruginosa* with inhibition zone diameters ranging from 13.67±7.37 to 17.00±2.65 mm and MIC values of 4-6µL/mL. *B. subtilis* and *S. aureus* were less sensitive (diameters of the zone of inhibition: 11.00±2.00 and 12.33±1.13 mm respectively, MIC: 8-10 µL/mL). On the other hand, the *R. lonadioides* EO showed weak activity against *L. monocytogenes*, *E. faecalis* and *E. coli* with diameters of inhibition zones ranging from 6.00±0.00 to 8.00±0.00 mm. According to these findings, it could be observed that the sensitivity of the microbial strains to our *R. lonadioides* essential oil was not correlated to the bacterial types (Gram-positive or Gram-negative). Hence, the studied essential oil could affect both of the bacterial membranes. Generally, the mechanisms of action of EOs and their selectivity towards certain bacteria have so far remained poorly understood (Hammer *et al.*, 1999, Bagamboula *et al.*, 2004). Accordingly, this selectivity is the result of the varied composition of the active fractions of the EOs, which often exhibit synergistic actions. It seems that their mechanisms of action are essentially linked to the structure of the wall and to the membrane permeability of Gram-positive and Gram-negative bacteria. In this context, the work of (Burt, 2004) showed that an active EO will exert its antimicrobial effect by its interference with the lipid bilayer of the target cell thanks to its hydrophobic property, which leads to a disturbance of the permeability and loss of the constituents of the cell. In addition, this reaction varies depending on the nature of the lipid bilayer, which explains the resistance of Gram-negative bacteria (Mahmoud *et al.* 2004). In addition, (Debbah *et al.*, 1970) demonstrated the high sensitivity of Gram-positive bacteria compared to Gram-negatives. In the same study approach, (Gordon *et al.*, 1973) and (Mahmoud *et al.*, 2004) suggested that the antimicrobial effect exerted by EOs could be explained by the destruction of certain enzymatic systems including those which participate in the production of cellular energy and the production of structural compounds. Moreover, (Mahmoud *et al.*, 2004), (Guesmi and Boudabous, 2006) for their part, put forward the hypothesis of inactivation and destruction of genetic material and,

finally (Caillet *et al.*, 2007) reported that EOs prevent the multiplication of bacteria, their sporulation and the synthesis of their toxins.

In the whole, as it was previously reported by many researchers that the antimicrobial activity of the EOs depends on their chemical composition, the obtained antimicrobial activity of our *R. lonadioides* essential oil towards the tested bacteria could be attributed to one or more of its major biological components: β-Pinene (41.85%), Phenanthroquinone (10.39%), D-Limonene (6.79%). However, α and β-Pinene have been reported to exhibit slight activity against a range of microorganisms with MIC values ranged from 7.5 to 20.0 mg/mL against *E. coli*, *S. aureus* and *E. faecalis* (Dorman and Deans, 2000; Sonboli *et al.*, 2006; Leite *et al.*, 2007; Jung, 2009; Runyoro *et al.*, 2010). Indeed, it is important to notice that synergistic interactions could occur between major or minor components of the EO thus playing a key effect on its antimicrobial activity (Dridi *et al.*, 2020).

By understanding the antibacterial activity of the Eos extracted from some plants of the *Anthemis* genus, the *Anthemis xylopoda* O. Schwarz EO (Borneol, 31.80%; β-pinene, 12.67%) showed a weak antibacterial effect against *B. subtilis*, *E. coli*, and *P. aeruginosa* with inhibition zone diameters ranging from 7.6 to 10 mm, while the antimicrobial activity against *S. aureus* was remarkably higher (23.2 mm) (Dridi *et al.*, 2020).

In parallel, *Anthemis stiparum* subsp. *Sabulicola* EO (with Germacrene D(11.13%) and t-Cadinol(11.01%) as major compounds), also showed low antimicrobial activity against the same bacterial strains with MIC values between 50 and 100 µL/mL (Šarac *et al.*, 2014).

In short, it could be summarized that oxygenated terpenes, alcohols, aldehydes and ketones are active but with different specificity and activity levels, which could be linked firstly to the functional group but also to hydrogen binding parameters (Panizzi *et al.*, 1993; Adam *et al.*, 1998).

**Table 1:** GC/SM Components of *Rhetinolepis lonadioides* Coss.essential oil from Algeria.

|    | Components <sup>a</sup> | RI lit <sup>b</sup> | RI <sup>c</sup> | R.T <sup>d</sup> | Area%        |
|----|-------------------------|---------------------|-----------------|------------------|--------------|
| 1  | styrene                 | 870                 | 890.94          | 6.09             | 0.11         |
| 2  | α-Pinene                | 930                 | 931.67          | 7.61             | 1.01         |
| 3  | α-Fenchene              | 959                 | 977.81          | 9.50             | 3.64         |
| 4  | β-Pinene                | 972                 | 980.00          | 10.93            | <b>41.85</b> |
| 5  | α-Terpinene             | 1017                | 1020.18         | 11.52            | 0.03         |
| 6  | D-Limonene              | 1018                | 1031.44         | 12.13            | <b>6.79</b>  |
| 7  | beta.-Ocimene           | 1041                | 1048.07         | 13.04            | 0.12         |
| 8  | γ-Terpinene             | 1047                | 1057.99         | 13.58            | 0.07         |
| 9  | Terpinolene             | 1080                | 1087.93         | 15.22            | 0.32         |
| 10 | Hotrienol               | 1115                | 1116.63         | 16.91            | 0.12         |
| 11 | .D-Camphor              | 1141                | 1140.76         | 18.42            | 0.22         |
| 12 | Borneol                 | 1148                | 1149.66         | 19.78            | 0.10         |
| 13 | α-Terpineol             | 1172                | 1188.40         | 21.39            | 0.25         |
| 14 | Bornylacetate           | 1285                | 1284.58         | 27.57            | 2.04         |
| 15 | Guaiacol<4-vinyl->      | 1293                | 1310.99         | 29.26            | 0.13         |
| 16 | Myrtenylacetate         | 1314                | 1324.01         | 30.08            | 0.32         |
| 17 | δ-Elmene                | 1334                | 1335.83         | 30.83            | 0.05         |
| 18 | Methyleugenol           | 1402                | 1400.65         | 35.15            | 0.82         |
| 19 | Caryophyllene           | 1421                | 1416.21         | 35.87            | 1.07         |
| 20 | α-Humulene              | 1454                | 1449.75         | 37.90            | 0.06         |
| 21 | (E)-β-Famesene          | 1448                | 1458.38         | 38.42            | 3.19         |

|                          |                                 |       |         |       |              |
|--------------------------|---------------------------------|-------|---------|-------|--------------|
| 22                       | $\beta$ -Selinene               | 1469  | 1467.02 | 38.95 | 1.61         |
| 23                       | $\alpha$ -Curcumene             | 1472  | 1478.90 | 39.67 | 1.63         |
| 24                       | $\alpha$ -Farnesene             | 1499  | 1508.29 | 41.43 | 0.24         |
| 25                       | $\delta$ -Cadinene              | 1514  | 1511.44 | 41.61 | 0.14         |
| 26                       | Nerolidol                       | 1605  | 1564.00 | 47.20 | 0.17         |
| 27                       | $\alpha$ -Bisabolol             | 1683  | 1681.11 | 51.19 | 0.14         |
| 28                       | Eudesm-7(11)-en-4-ol            | 1681  | 1686.59 | 51.49 | 0.47         |
| 29                       | Ethanone, 1-[1,1'-biphenyl]-4-y | 1716  | 1739.56 | 54.30 | 2.66         |
| 30                       | NI                              | ----- | 1793.37 | 57.12 | 5.26         |
| 31                       | NI                              | ----- | 1817.50 | 58.34 | <b>6.93</b>  |
| 32                       | Phenanthroquinone               | 1880  | 1867.70 | 60.85 | <b>10.39</b> |
| 33                       | NI                              | ----- | 1890.70 | 62.00 | 0.52         |
| 34                       | Farnesylacetone                 | 1895  | 1897.07 | 62.32 | 0.24         |
| 35                       | m-Camphorene                    | 1944  | 1954.24 | 65.05 | <b>5.43</b>  |
| 36                       | $\alpha$ -Springene             | 1969  | 1969.42 | 66.59 | 1.85         |
| Hydrocarbonmonoterpenes  |                                 |       |         |       | 53.94%       |
| Oxygenatedmonoterpenes   |                                 |       |         |       | 14.39%       |
| Hydrocarbosesquiterpenes |                                 |       |         |       | 7.99%        |
| Oxygenatedsesquiterpenes |                                 |       |         |       | 3.68%        |
| Hydrocarbonditerpenes    |                                 |       |         |       | 7.28%        |
| Total identified (%)     |                                 |       |         |       | 80.00%       |

**Notes:**

<sup>a</sup>Order of elution and percentages of individual components are given on an apolar column (DB-5).

<sup>b</sup>RI lit, retention indices taken in the literature, ref. (60 ; 61), <sup>c</sup>RI: Retention index calculated from relative retention times, <sup>d</sup>RT : retention times.

**Table 2:** Antimicrobial activity of *Rhetinolepis lonadioides* essential oil (aerial parts): Inhibition zones and MICs.

| Bacteria                      | Inhibition zones (mm)             |   |  | MIC of <i>R. lonadioides</i> essential oil ( $\mu\text{L}/\text{mL}$ ) |
|-------------------------------|-----------------------------------|---|--|--|
|                               | Essential oil (15 $\mu\text{L}$ ) | Spiramycin (at 100 $\mu\text{g}/\text{d}$ ) | Oflatoxin (at 5 $\mu\text{g}/\text{d}$ ) |  |
| <b>Gram-positive bacteria</b> |                                   |   |  |  |
| <i>Bacillus cereus</i>        | 17.00 $\pm$ 2.65                  | 17.67 $\pm$ 2.08                            | 22.00 $\pm$ 1.73                         | 4  |
| <i>Bacillus subtilis</i>      | 11.00 $\pm$ 2.00                  | 16.67 $\pm$ 1.53                            | 20.67 $\pm$ 1.15                         | 10   |
| <i>Staphylococcus aureus</i>  | 12.33 $\pm$ 1.13                  | 17.33 $\pm$ 1.15                            | 24.00 $\pm$ 1.00                         | 8  |
| <i>Enterococcus faecalis</i>  | 8.00 $\pm$ 1.5                    | 22.67 $\pm$ 2.08                            | 26.67 $\pm$ 1.53                         | --   |
| <b>Gram-negative bacteria</b> |                                   |   |  |  |
| <i>Listeria monocytogenes</i> | 6.00 $\pm$ 0.00                   | 30.20 $\pm$ 2.00                            | 21.67 $\pm$ 2.08                         | --   |
| <i>Escherichia coli</i>       | 6.00 $\pm$ 0.00                   | 6.00 $\pm$ 0.00                             | 41.00 $\pm$ 5.57                         | --   |
| <i>Pseudomonas aeruginosa</i> | 13.67 $\pm$ 7.37                  | 20.67 $\pm$ 5.03                            | 26.67 $\pm$ 3.06                         | 6  |

**Conclusion**

The present study investigated, for the first time, the chemical composition of *R. lonadioides* essential oil endemic of Algeria. The studied essential oil is dominated by monoterpene hydrocarbons (53.94%), oxygenated monoterpenes (14.39%), oxygenated sesquiterpenes (3.68%) and sesquiterpene hydrocarbons (7.99%). The main compounds are  $\beta$ -Pinene (41.85%), Phenanthroquinone (10.39%) and D-Limonene (6.79%). Furthermore, our results showed that the essential oil of *R. lonadioides* possessed interesting antimicrobial properties which potentially suggest its exploitation as a new natural antimicrobial agent in pharmaceutical applications for treating or preventing human infectious diseases and/or in food industries.

**Acknowledgements**

Authors would like to thank Professor A. Marouf (University of Nâama, Algeria) for the identification of the plant material.

**References**

Adam, K.; Sivropoulou, A.; Kokkini, S.; Lanaras, T. and Arsenakis, M. (1998). Antifungal activities of

*Origanum vulgare* subsp. *hirtum*, *Menthaspicata*, *Lavandula angustifolia* and *Salvia fruticosa* essential oils against human pathogenic fungi. *J. Agricultural Food Chem.*; 46: 1739-1745.

African Plant Database (2021). Conservatoire et Jardin botaniques de la Ville de Genève and South African National Biodiversity Institute, Pretoria. Available from: <http://www.ville-ge.ch/musinfo/bd/cjb/africa/recherche.php> (Retrieved June 11, 2021).

Aghajani, Z.; Masoudi, S. and Rustaiyan, A. (2005). Volative oils of *Anthemis talyshensis* A. Fedor and *Sclerorhachis platyrachis* (Boiss.) Podlech ex Rech.f from Iran. *JEssent. Oil Res.*; 17: 355-357.

Albayrak, S. and Aksoy, A. (2012) Evaluation of antioxidant and antimicrobial activities of two endemic *Anthemis* species in Turkey. *J. Food Biochem.*; 37(6): 639-645.

Alessandro, V.; Claudio, F.; Giacomo, R.; Mirella, D.C.; Giampiero, C.; Mauro, S. and Armandodoriano, B. (2016). Secondary metabolites with ecologic and medicinal implications in *Anthemis cretica* subsp. *petraea* from Majella National Park. *AIMS Mol. Sci.*; 3(4): 648-660.

- Bagamboula, C.F.; Uyttendaele, M.; Candan, F.; Daferera, D.; Unli, G.V.; Polissiou, M. and Sokmen, A. (2004). Antimicrobial and antioxidative activities of the essential oils and methanol extracts of *Salvia crytanha* (Monbret et Aucher ex Benth and *Salvia multicaulis* (Vahl.) *Food Chem.*; 84: 519-525.
- Battandier, J. (1888). Flore de l'Algérie. Ancienne flore d'Alger transformée contenant la description de toutes les plantes signalées jusqu'à ce jour comme spontanées en Algérie : Dicotylédones. A. Jourdan, Alger / Paris.
- Baytop, T. (1999). Türkiye'de bitkiler ile tedavi, geçmiştene bugün (Therapy with Medicinal Plants in Turkey, Past and Present), 2nd ed. Nobel Tıp Kitabevleri, Istanbul, Turkey.
- Bruno, M.; Bono, M.L.; Vassallo, N.; Gedris, T.E. and Herz, W. (1997). Guaianolides and other terpenoids from *Anthemis aetnensis*. *Phyto. Chem.*; 45: 375-377.
- Bulatovic, V.M.; Menkovic, N.R.; Vajs V.E.; Milosavljevic, S.M. and Djokovic, D.D. (1998). Essential oil of *Anthemis Montana*. *J. Essent. Oil Res.*; 10: 223-226.
- Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods-a review. *Int. J. Food Microbiol.*; 94: 223-253.
- Caillet, S. and Lacroix, M. (2007). Les huiles essentielles : leurs propriétés antimicrobiennes et leurs applications potentielles en alimentaire. Laboratoire de Recherche en Sciences Appliquées à l'Alimentation (RESALA) INRS-Institut Armand-Frappier, Université de Laval (Québec).
- Chemsa, A.E.; Zellagui, A.; Öztürk, M.; Erol, E.; Ceylan, O.; Mehmet Emin Duru, M.E.M. Lahouel (2018). Chemical composition, antioxidant, anticholin esterase, antimicrobial and antibiofilm activities of essential oil and methanolic extract of *Anthemis stiparum* subsp. *Sabulicola* (Pomel) *Oberpr. Microb. Pathog.* 119: 233-240.
- Collu, F.; Bonsignore, L.; Casu, M.; Floris, C.; Gertsch, J. and Cottiglia, F. (2008). New cytotoxic saturated and unsaturated cyclohexanones from *Anthemis maritima*; *Bioorg. Med. Chem. Lett.* 18: 1559-1562.
- Dabbah, R.; Edwards, V.M. and Moats, W.A. (1970). Antimicrobial action of some citrus fruit oils on selected food-borne bacteria. *Appl. Microbiol.*; 19: 27-30.
- Dorman, H.J. and Deans, S.G. (2000). Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J. Appl. Microbiol.*; 88: 308-316.
- Dridi, I.; Haouel-Hamdi, S.; Cheraief, I.; Mediouni Ben Jemâa, J.; Landoulsi, A. and Chaouch, R. (2020). Essential oil from flowering tops of *Lavandula dentata* (L.). Chemical composition, antimicrobial, antioxidant and insecticidal activities. *J. Essent. oil Bear. Plants*, 24: 632-647.
- Gonenc, T.; Argyropoulou, C.; Erdogan, T.; Gousiadou, C.; Juergenliemk, G.; Kivcak, B. and Skaltsa, H. (2011). Chemical constituents from *Anthemis wiedemanniana* Fisch. & Mey. *Biochem. Syst. Ecol.*, 39: 51-55.
- Gordon, R.E.; Haynes, W.C. and Pang, C.H.N. (1973) The genus *Bacillus*. *Agriculture Handbook*. N° 427, ARS-USDA, Washington (USA).
- Grace, M.H. (2002). Chemical composition and biological activity of the volatiles of *Anthemis melampodina* and *Pluchea dioscoridis*. *Phytother. Res.* 16: 183-185.
- Guesmi, A. and Boudabous, A. (2006). Activité antimicrobienne de cinq huiles essentielles associées dans les produits de thalassothérapie- les Plantes à Parfum, Faculté des sciences de Tunis, université Tunis-El Manar, Tunisie.
- Guragaç Dereli F.T.; Ilhan, M. and EsraKüpeliAkkol E.K. (2018). Discovery of new antidepressant agents: *In vivo* study on *Anthemis wiedemanniana* Fisch. & Mey. *J. Ethnopharmacol.* 226: 11-16.
- Hammer, K.A.; Carson, C.F. and Riley, T.V. (1999). Antimicrobial activity of essential oils and other plant extracts. *J. Appl. Microbiol.*, 86: 985-990.
- Hofer, O. and Greger, H. (1985) New sesquiterpene-coumarin ethers from *Anthemis cretica*. *Liebigs Annalen der Chemie*, 6: 1136-1144.
- Holla, M.; Svajdlenka, E.; Vaverkova, S.; Zibrunova, B.; Tekel, J. and Havranek, E. (2002). Composition of the oil from flower heads of *Anthemis tinctoria* L. cultivated in Slovak republic. *J. Essent. Oil Res.*, 12: 714-716.
- Jalali-Heravi, M.; Moazeni-Pourasil, R.S. and Sereshti, H. (2015). Elimination of chromatographic and mass spectrometric problems in GC-MS analysis of lavender essential oil by multivariate curve resolution techniques: improving the peak purity assessment by variable size moving window evolving factor analysis. *J. Chromatogr B Analyt. Technol. Biomed. Life Sci.* 83(9): 983-984.
- Javidnia, K.; Miri, R.; Kamalinejad, M.; Sarkarzadeh, H. and Jamalian, A. (2004). Chemical composition of the essential oils of *Anthemis altissima* L. grown in Iran. *Flavour Fragr. J.* 19: 213-216.
- Jung, E.K. (2009). Chemical composition and antimicrobial activity of the essential oil of *Chrysanthemum indicum* against oral bacteria. *J. Bacteriol. Virol.*, 39: 61-69.
- Kivcak, B.; Mert, T.; Saglam, H.; Ozturk, T.; Kurkcuglu, M. and Baser, K.H. (2007). Chemical composition and antimicrobial activity of the essential oil of *Anthemis wiedemanniana* from Turkey. *Chem. Nat. Compd.*, 43: 47-51.
- Korkmaz, M. and Karakus, S. (2015). Traditional uses of medicinal plants of Üzümlü district, Erzincan, Turkey. *Pak. J. Bot.*, 47: 125-134.
- Kultur, S. (2007). Medicinal plants used in Kirklareli Province (Turkey). *J. Ethnopharmacol.* 111: 341-364.
- Kürkçüoğlu, M. and Tosun, F. (2020). Determination of the volatile compounds of *Anthemis cretica* sub sp. *anatolica* (Boiss.) Grierson *Nat. Nat. Volatiles Essent. Oils* 7(2): 10-16.
- Kurtulmus, A.; Fafal, T.; Mert, T.; Saglam, H.; Kivcak, B.; Ozturk, T.; Demirci, B. and Baser, K. (2009). Chemical composition and antimicrobial activity of the essential oils of three *Anthemis* species from Turkey. *Chem. Nat. Compd.*, 45(6): 900-904.
- Leite, A.M.; Lima, O.; Souza, E.L.; Diniz, F.F.; Trajano, V.N. and Medeiros, I.A. (2007). Inhibitory effect of  $\beta$ -pinene,  $\alpha$ -pinene and eugenol on the growth of potential infectious endocarditis causing Gram-positive bacteria. *Revista Brasileira de Ciências Farmacêuticas*, 43: 121-126.
- Mahmoud, B.; Yamazaki, K.; Miyashita, K.; IL-Shik, S.; Dong-Suk, C. and Suzuki, T. (2004). Bacterial microflora of carp (*Cyprinus carpio*) and its shelf-life

- extension by essential oil compounds. *Food Microbiol.*, 21: 657-666.
- NCCLS (National Committee for Clinical Laboratory Standards) (1997) Performance standards for antimicrobial disk susceptibility test, 6th edn. Approved Standard M2-A6, Wayne, PA.
- Oberprieler, C.H. (1998). The systematics of *Anthemis* L. (*Compositae*, *Anthemideae*) in W and C North Africa. *Boccone* 9: 1-328.
- Oberprieler, C.H.; Vogt, R. and Watson, L.E. (2007). Tribe *Anthemideae* Cass. The Families and Genera of Vascular Plants. In: Kubitzki K (editor). Flowering Plants, Eudicots, Asterales, Berlin, Germany: Springer 8: 342-374.
- Orlando, G.; Zengin, G.; Ferrante, C.; Ronci, M.; Recinella, L.; Senkardes, I.; Gevrenova, R.; Zheleva-Dimitrova, D.; Chiavaroli, A.; Leone, S.; Di Simone, S.; Brunetti, L.; Picot-Allain, C.M.N.; Mahomoodally, M.F.; Sinan, K.I. and Menghini, L. (2019). Comprehensive chemical profiling and multidirectional biological investigation of two wild *Anthemis* species (*Anthemis tinctoria* var. *pallida* and *A. cretica* subsp. *tenuiloba*): focus on neuroprotective Effects. *Mol.* 24: 25-82.
- Panizzi, L.; Flamini, G.; Cioni, P.L. and Morelli, I. (1993). Composition and antimicrobial properties of essential oils of four Mediterranean *Lamiaceae*. *J. Ethnopharmacol.* 39: 167-170.
- Pavlovic, M.; Lakusic, D.; Kovacevic, N.; Tzakou, O. and Couladis, M. (2010). Comparative analysis of essential oils of six *Anthemis* taxa from Serbia and Montenegro. *Chem. Biodivers.* 7: 1231-1244.
- Rates, S.M. (2001). Plants as source of drugs. *Toxicol.* 39: 603-613.
- Runyoro, D.; Ngassapa, O.; Vagionas, K.; Aliyannis, N.; Graikou, K. and Chinou, I. (2010). Chemical composition and antimicrobial activity of the essential oils of four *Ocimum* species growing in Tanzania. *Food Chem.* 119: 311-316.
- Rustaiyan, A.; Azar, P-A.; Moradalizadeh, M.; Masoudi, S. and Ameri, N. (2004). Volatile constituents of three Compositae herbs: *Anthemis altissima* L. var. *Altissima*, *Conyza canadensis* (L.) Cronq. and *Grantinaaicheri* Boiss. growing wild in Iran, *J. Essent. Oil Res.*, 16: 579-581.
- Sadiki, F.Z. and El-Idrissi, M. (2019). Chemical composition of essential oil of *Anthemis nobilis* L. flowers from Morocco. *Appl. J. Envir. Eng. Sci.*, 5(4): 342-348.
- Šarac, Z.; Matejić, J.S.; Stojanović-Radić, Z.Z.; Veselinović, J.B.; Džamić, A.M.; Bojović, S. and Marin, P.D. (2014). Biological activity of *Pinus nigra* terpenes-evaluation of Fts Z inhibition by selected compounds as contribution to their antimicrobial activity. *Comput. Biol. Med.*, 54: 72-78.
- Saroglou, V.; Dorizas, N.; Kypriotakis, Z. and Skaltsa, H.D. (2005). Analysis of the essential oil composition of eight *Anthemis* species from Greece. *J. Chromatography A* 11(4): 313-322.
- Sonboli, A.; Babakhani, B. and Mehrabian, A. (2006). Antimicrobial activity of six constituents of essential oil from *Salvia*. *Zeitschrift für Natur for schung*, 61: 160-164.
- Stojkovic, N.; Stojkovic, M.; Marinkovic, M.; Chopra, G.; Kostic, D. and Zarubica, A. (2014). Polyphenol content and antioxidant activity of *Anthemis cretica* L. (Asteraceae). *Oxi. Commun.* 37(1): 237-246.
- Tawaha, K.A.; Alali, F.Q. and Hudaib, M.M. (2015). Chemical composition and general cytotoxicity evaluation of essential oil from the flowers of *Anthemis palestina* Reut. ex Boiss.; growing in Jordan. *J. Essent. Oil Bear. Plants* 18(5): 1070-1077.
- Toiu, A.; Muntean E.; Oniga I. and Tămaș, M. (2009). Pharmacognostic Research on *Viola declinata* Waldst. et Kit (Violaceae), *Farmacia*, 57(2): 218-22.
- Ugurlu, E. and Secmen, O. (2008). Medicinal plants popularly used in the villages of Yunt Mountain (Manisa-Turkey). *Fitoterapia*, 79: 126-131.
- Uzel, A.; Guvensen, A. and Cetin, E. (2004). Chemical composition and antimicrobial activity of the essential oils of *Anthemis xylopoda* O. Schwarz from Turkey. *J. Ethnopharmacol.* 95: 151-154.
- Vajs, V.; Bulatovic, V.; Fodulovic-Savikin, K.; Menkovic, N.; Macura, S.; Juranic, N. and Milosavljevic, S. (1999). Highly oxygenated guaianolides from *Anthemis cretica* subsp. *cretica*. *Phytochem.*, 50: 287-291.
- Venditti, A.; Frezza, C.; Rossi, G.; Di Cecco, M.; Ciaschetti, G.; Serafini, M. and Bianco, A. (2016). Secondary metabolites with ecologic and medicinal implications in *Anthemis cretica* subsp. *Petraea* from Majella National Park. *AIMS Mol. Sci.*, 3: 448-660.
- Yusufoglu, H.S.; Tabanca, N.; Bernier, U.R.; Li, A.Y.; Salkini, M.A.; Alqasoumi, S.I. and Demirci, B. (2018). Mosquito and tick repellency of two *Anthemis* essential oils from Saudi Arabia. *Saudi Pharmaceutical J.*, 26: 860-864.