



Plant Archives

Journal home page: www.plantarchives.org

DOI Url: <https://doi.org/10.51470/PLANTARCHIVES.2021.v21.no1.246>

STUDIES ON THE EFFECT OF ARBUSCULAR MYCORRHIZA AND PHOSPHATE SOLUBILIZING BACTERIA UNDER REDUCED FERTILIZER APPLICATION ON TOMATO

K. Sivakumar and N. Pandeewari

Department of Microbiology, Faculty of Agriculture, Annamalai University, Tamil Nadu., India

(Date of Receiving-05-01-2021; Date of Acceptance-03-04-2021)

ABSTRACT

A field experiment was carried out at department of agricultural microbiology, Annamalai University, Chidambaram, Tamil Nadu, India to evaluate the performance of tomato inoculated with Arbuscular Mycorrhiza (AM) and phosphate solubilizing bacteria (PSB) when the fertilizer application was reduced to 50% and 25% of the recommended fertilizer dose (RDF). The results of the study revealed that, inoculation of tomato with AM and PSB significantly increased all the parameters examined during the study. However, the best results were obtained when the seedlings were inoculated with AM followed by PSB at 50% of the RDF as evident from 16.6% and 12.6% increase in the total plant dry weight at the respective treatments as compared to control. Similarly the concentration of nitrogen (N), phosphorus (P), and potassium (K) in the plants inoculated with AM at 50% of the RDF was increased by 17.3, 20.6 and 23.2% followed by an increase of 2.5, 11.5, and 12.2% at PSB with 50% RDF as compared to control. Yield of tomato was respectively increased by 9.7% and 4.4% when inoculated with AM and PSB at 50% of RDF as compared to control.

Keywords: Tomato, fertilizer, inoculation, PSB and AM

INTRODUCTION

Tomato (*Lycopersicon esculentum* L.) is one of the most studied fleshy fruits, and it has assumed the status of “functional food” because of the epidemiological evidence as regards the reduced risk of cancer and cardiovascular diseases in relation to its consumption. Tomato fruits contain high concentrations of antioxidant molecules such as lycopene, ascorbic acid and carotenoids (Giovannetti *et al.*, 2012). Tomato (*Lycopersicon esculentum* L.) is the second most important fruit or vegetable crop next to potato (*Solanum tuberosum* L.), with approximately 182.3 million tons of tomato fruits produced on 4.85 million ha each year (FAOSTAT, 2019).

AM fungi is the most abundant kind of mycorrhizae found in association with every taxonomic group of plants and the list of species not infected is probably far shorter than the infected ones. These fungal associations are beneficial to crop plants in many ways, including enhancing the nutrient availability especially phosphorus, enhancing water uptake, inducing resistant against diseases and increasing the yield (Lekberg and Koids, 2005).

Phosphorus solubilizing biofertilizers are carrier based preparations containing living cells of microorganisms like bacteria, fungi and actinomycetes which may help in increasing crop productivity by way of helping in solubilization of insoluble phosphorus, stimulating plant growth by providing hormones, vitamins and other growth promoting substances.

Phosphate Solubilizing Bacteria (PSB) are capable of

hydrolyzing organic and inorganic phosphorus from insoluble compounds and PSB produce phosphatase like phytase that hydrolyse organic forms of phosphate compounds efficiently (Zehra, 2010).

The chemical fertilizers like N, P and K have played a significant role in increasing the yield and quality of plant products during early seventies. However, indiscriminate use of chemical fertilizers has resulted in several problems like loss of fertility and multiple nutrient deficiencies. In addition to it, heavy inoculation of inorganic fertilizers degrade the soil health by adversely affecting the microbial biodiversity, physical and chemical environment of soil, water bodies and capital inputs like soil, water and thus overall ecology which ultimately result in reduced crop productivity and quality (Ghanti and Sharangi, 2009). The reason for the increased use of chemical fertilizers is the increased demand for food and the ability of some of the mineral elements to get fixed with soil colloids after application. For example P is an essential macro element for plants, yet the total concentration of P in soils ranges from 0.02- 0.5%. Thus, to increase the availability of phosphorus for plants, large amount of fertilizers are used on a regular basis, because after inoculation, a large proportion of phosphorus is quickly transferred to an insoluble form (Omar, 1997).

The application of biofertilizers has a significant effect on growth and yield of different vegetables. However, information on synergism between the different biofertilizers under reduced fertilizer application is still lacking. In this perspective it was thought worthwhile to carry out an experiment to find out the effect of AM

fungus and PSB on different morphological, quality and yield parameters of under reduced inorganic fertilizer application.

MATERIAL AND METHODS

A field experiment was carried out on department of Agricultural Microbiology, Annamalai University, Chidambaram, Tamil Nadu in India. The soil of the experimental field is sandy loam with an EC of 0.33 $\mu\text{S}/\text{cm}$ and pH 7.3.

Land preparation and fertilizer application

The experimental area was thoroughly prepared by ploughing three times and well decomposed farm yard manure (FYM) was applied at the rate of 25 t/ha followed by the preparation of plots. Fertilizers were applied to the plots as per the treatments as shown in table-1. N was applied to the plots in two doses, one at the time of transplanting and second in the month of June, whereas the entire P and K was applied in a single dose at the time of transplanting as per the designed treatments.

Treatments

Seven treatments shown in the table-1 were employed which were laid out in a randomized block design and replicated thrice. AM was applied in the root zone at the time of transplanting @ 10 Kg/ha, whereas seedlings were dipped in a solution of PSB for 45 minutes before transplanting.

Table 1: Treatment details

Symbol	Treatment
T0	Control
T1	50 % RDF
T2	25 % RDF
T3	50 % RDF + AM
T4	50 % RDF + PSB
T5	25 % RDF + AM
T6	25 % RDF + PSB

Intercultural operations

All the required cultural practices such as irrigation, weeding, pest and disease control etc. were kept constant and given uniformly in all the experimental plots. Ten plants were randomly selected from each plot in such a way that the marginal effect was avoided and data were recorded on the following parameters at the time of harvest.

i. **Plant height:** Plant height was measured with the help of a scale.

ii. **Plant dry weight:** For calculation of plant dry weight, samples were oven dried at 80 °C till their weight became constant.

iii. **Chlorophyll content:** Chlorophyll content was estimated at the time of second picking i.e. 85 DAT. The procedure developed by Witham *et al.*, (1971) was followed for chlorophyll estimation.

iv. **Nitrogen (N):** N content from the bulbs was determined by using the method as proposed by Peoples *et al.*, (1989)

v. **Phosphorus (P):** P content was estimated from the bulbs as per the method proposed by Koenig and Johnson (1942).

vi. **Potassium (K):** K was estimated by flame photometric method as proposed by Reed and Scott (1962).

vii. **Total soluble solids (TSS):** TSS of the bulbs was measured by using the hand refractometer.

viii. **Yield:** The ripe fruit yield obtained from each plot was recorded in all the pickings. The yield was calculated and expressed in tons per hectare.

Data reported in this study are the mean values based on the three replicates. Differences among treatments were tested by ANOVA and the mean values among the treatments were compared using Duncan's multiple range test at $p=0.05$.

RESULTS AND DISCUSSION

Plant height and total dry weight

Plant height observed a continuous decrease when the fertilizer application was reduced to 50 % and 25 % of the RDF. However, the inoculation with AM and PSB significantly increased the plant height as compared to control and the non inoculated plants which received a reduced fertilizer dose (table-2). Data indicates that the maximum increase in the plant height was observed in the plants inoculated with AM followed by PSB at 50 % of the RDF. Height of the plants at these treatments respectively observed an increase of 8.4% and 1.9 % increase as compared to control. Similar to plant height, total plant dry weight also followed a decreasing trend with reduction in the fertilizer application to 50 % and 25 % of the RDF. Total plant dry weight was significantly increased with inoculation of AM and PSB as evident from table-2. However, maximum increase of 16.6 % was observed by inoculating the plants with AM at 50 % of the RDF followed by an increase of 12.6 % in the plants treated with PSB at the same level of the fertilizer dose as compared to control. The results are in agreement with Dar *et al.*, (2018) and Naik (2014) who reported an increase in the plant height of onion through inoculation with AM

and PSB at reduced phosphorus application which can be because of higher solubilization of P brought about by PSB followed by its increased uptake along with N and K as is evident from table- 2 of this study. Inoculation with AM increases the absorbing capacity of the roots favoring more water uptake by the plants resulting in cell expansion responsible for the increased plant height (Huixing Song, 2005). It has been found that AM fungus improves the hydraulic conductivity and water uptake of roots hence providing a low resistance pathway for radial flow of water across cortex, and contributing towards better water uptake by the plants (Kothari *et al.*, 1990). Furthermore, AM fungus may also regulate the hydraulic properties of plants through regulation of plasma membrane intrinsic proteins in combination with phytohormones (Ruiz-Lozano *et al.*, 2009). Since nitrogen is a constituent of chlorophyll, the increase uptake of which might have resulted in increased synthesis of photosynthetic leading to better plant growth. The second major nutrient phosphorus is an essential constituent of cellular protein and nucleic acid which might have encouraged the cell division and meristematic activity of plants resulting in increased plant height and total plant dry weight (Monika *et al.*, 2006).

Chlorophyll content

Reduction in fertilizer application to 50 % and 25 % of the RDF had an adverse affect on the plant chlorophyll content (table-2). However, inoculation with AM and PSB led to increase in the chlorophyll content at all the treatments. However, the maximum increase was observed in the plants subjected to inoculation with AM followed by PSB at 50 % of the recommended fertilizer dose. Chlorophyll content at these treatments was respectively increased by 12.1% and 9.2% as compared to control.

The AM inoculated plants have shown to have higher chlorophyll content compared to non AM plants. This observation has been made by several workers in different systems (Auge, 2001) and Colla *et al.*, (2008). In pepper for instance, the chlorophyll content was shown to be increased by 15-25% to indicate the relevance of mycorrhizal association in increasing the pigment content in the leaves (Beltrano *et al.*, 2013). It has been well recognized that chlorophyll concentration is related to photosynthetic rate and chlorophyll fluorescence. Thus, in AM inoculated plants an increased rate of chlorophyll has been associated with the increased rate of photosynthesis or with higher magnesium and nitrogen which are major constituents of chlorophyll (Mathur and Vyas, 1995).

Effect on mineral uptake

Mineral uptake observed a continuous decrease with the reduction in fertilizer application by 50 % and 25 % of RDF. However, it was significantly improved with the inoculation of plants by AM and PSB. It is evident from

table-3 that N content in the plant tissues was decreased by 55.2 and 68.4 % as compared to control when the RDF was reduced respectively by 50 % and 25 %. P concentration at 50 % and 25 % of the RDF was decreased by 17.3 and 60.8 percent as compared to control followed by a reduction of 27.9 and 67.4% in K concentration in the plant tissues at the same treatments. Table-3 clearly indicates an increase in the concentration of N, P and K in the plants that were inoculated with AM and PSB. However the maximum increase was obtained in the AM inoculated plants treated with 50 % of the RDF. N, P and K in the plant tissues treated with 50 % of the RDF was increased by 17.3, 20.6 and 23.2 % as compared to control followed by an increase of 2.5, 11.5 and 12.2 % in the plants subjected to inoculation with PSB at 50 % of the RDF. Similar results have also been found by Talwar *et al.*, (2016) and Hashem (2015). Microorganisms can increase the solubility of inorganic P by releasing protons, H⁺ and organic acid anions such as citrate, malate and oxalate (Yousefi *et al.*, 2011).

whereas the AM fungus absorbs the mineral elements like N, P, K, Fe, Zn etc much faster through physical exploration of the soil, modification of root environment, increased movement of various mineral elements into mycorrhizal fungus hyphae and their translocation to the associated plant (Turk *et al.*, 2006). AM fungi expand the roots by adding their own expansive network of absorbing strands to mine the soil for water and the dissolved minerals carried therein may also alter the water relations and the drought tolerance of the plant. It has been found that AM fungus improves the hydraulic conductivity and water uptake of roots hence providing a low resistance pathway for radial flow of water across cortex, and contributing towards better water uptake by the plants (Kothari *et al.*, 1990) [12]. Furthermore, AM fungus may also regulate the hydraulic properties of plants through regulation of plasma membrane intrinsic proteins in combination with phytohormones (Ruiz-Lozano *et al.*, 2009). Tinker (1984) while summarizing the effects of AM on increased water uptake indicated that the probable reason for increased water uptake are as the better distribution of absorbing hyphal network, greater surface area and better extension rate, altered soil rhizosphere properties, uptake kinetics, faster water extraction from reduced water potential soils and hence faster recovery from the water stress.

Yield

The yield was severely reduced in tomato due to reduction in fertilizer application. The yield was reduced by 26.1 % and 50.7 % when the fertilizer application was reduced to 50 % and 25 % of the RDF. Inoculation with AM and PSB significantly increased the yield even at the reduced fertilizer levels. However, AM fungus was found to have highest effect on the yield of tomato at 50 % of the RDF where the yield was increased by 9.7% followed by an increase of 4.4 % at the same fertilizer dose and PSB

inoculation as compared to control. This effect was in conformity with the results obtained by Subbiah (1994) in chilli and onion, Nanthakumar and Veeraragavathatham (2000) in brinjal and Yadav *et al.*, (2005) in onion. The significant increase in yield due to biofertilizers may be an account of its direct role in nitrogen fixation, production of phytohormone like substances and increased uptake of nutrients such as nitrogen. The increase in yield by the AM could be due to the function of mycorrhiza to enhance the efficiency of nutrient absorption and release of growth substances. This phenomenon of AM action might have increased the yield attributing characters in tomato as has been found by several workers *viz.*, Bagyaraj and Sreeramulu (1982) in chillies, and Subbiah (1994) in chilli and onion.

Table 2: Effect of reduced fertilizer application, arbuscular mycorrhiza and PSB on growth parameters and chlorophyll content of tomato

Treatments	Plant height (cm)	Plant dry weight (g)	Chlorophyll (mg g ⁻¹ FW)
Control	84.3	41.0	0.87
50 % RDF	63.2	27.4	0.46
25 % RDF	44.5	21.2	0.23
1/2RDF + VAM	91.2	47.2	0.99
1/2RDF + PSB	85.8	45.5	0.92
1/4RDF + VAM	60.3	35.2	0.65
1/4RDF + PSB	52.4	33.2	0.53
CD (5%)	4.4	1.2	0.03

Table 3: Effect of reduced fertilizer application, arbuscular mycorrhiza and PSB on quality parameters and yield of tomato

Treatments	Nitrogen (%)	Phosphorus (%)	Potassium (%)	Yield (t/ha)
Control	3.8	0.23	4.3	6.5
50 % RDF	1.7	0.19	3.1	4.8
25 % RDF	1.2	0.09	1.4	3.2
1/2RDF +VAM	4.6	0.29	5.6	7.2
1/2RDF +PSB	3.9	0.26	4.9	6.8
1/4RDF +VAM	2.3	0.10	2.3	4.2
1/4RDF +PSB	1.9	0.15	1.9	3.7
CD (5%)	0.4	0.04	0.3	0.35

REFERENCES

Auge RM. Water relations, drought and vesicular- arbuscular mycorrhizal symbiosis. *Mycorrhiza*. 2001; 11:3-42.
 Bagyaraja DJ, Sreeramulu KR. Pre-inoculation with VA

mycorrhiza improves growth and yield of chilli transplanted in the field saves phosphatic fertilizer. *Plant Soil*. 1982; 69:375-381.

- Beltrano J, Ruscitti M, Arango MC, Ronco M. Effects of arbuscular mycorrhiza inoculation on plant growth, biological and physiological parameters and mineral nutrition in pepper grown under different salinity and P levels. *Journal of Soil Science and Plant Nutrition*. 2013; 13:123-141.
- Colla G, Roupheal Y, Cardarelli M, Temperini O, Rea E, Salerno A, Pierandrei F. Influence of grafting on yield and fruit quality of pepper (*Capsicum annuum* L.) grown under greenhouse conditions *Acta Horticulture*. 2008; 782:359-363
- Ghanti S, Sharangi AB. Effect of bio-fertilizers on growth, yield and quality of onion cv. Sukhsagar. *Journal of Crop and Weed*. 2009; 5(1):120-123.
- Giovannetti, M. and L. Avio. 1985. VAM infection and reproduction as influenced by different organic and inorganic substances. In: Proceedings of 6th NACOM, (ed.), Molina, R., Foresh Research Laboratory, Oregon State University, Corvallis, p. 95.
- Hashem MM. Impact of Inoculation with arbuscular mycorrhizal, phosphate solubilizing bacteria and soil yeast on growth, yield and phosphorus content of onion plants. *International Journal of Soil Sciences*. 2015; 10(2):93-99.
- Huixing S. Effects of VAM on host plant in the condition of drought stress and its mechanisms. *Electronic Journal of Biology*. 2005; (3):44-48.
- Koenig R, Johnson C. Colorimetric determination of phosphorus in biological materials. *Industrial and Engineering Chemistry Analytical Edition*. 1942; 14(2):155-156.
- Kothari SK, Marschner H, George E. Effect of VA mycorrhizal fungi and rhizosphere microorganisms on root and shoot morphology, growth, and water relations of maize. *New Phytologist*. 1990; 116:303-
- Lekberg, Y. and R.T. Koids. 2005. Arbuscular mycorrhizal fungi, rhizobia available P and nodulation of groundnut (*Arachis hypogea* L.) in zimbabwe. *Agric. Ecosys. Environ.*, 110: 143-148.
- Monika K, Fernando AL, Agustin AG, Hans S. Phosphorus deficiency decreases cell division and elongation in grass leaves. *Plant Physiology*. 2006; 141(2):766-775.
- Naik MR. Effect of inorganic fertilizers, PSB and VAM on growth and yield of onion. *Mysore Journal of Agricultural Sciences*. 2014; 48(3):460-463.
- Nanthakumar S, Veeraragavathatham D. Effect of integrated nutrient management on growth parameters and yield of brinjal (*Solanum melongena* L) cv. PLR-1. *South Indian Horticulture*. 2000; 48(1-6):31-35.
- Omar SA. The role of rock-phosphate-solubilizing fungi and Vesicular-Arbuscular-Mycorrhiza (VAM) in growth of wheat plants fertilized with rock phosphate. *World*

- Journal of Microbiology and Biotechnology*. 1997;
- Peoples MB, Faizah AW, Rerkasem B, Herridge DF. Methods for evaluating nitrogen fixation by nodulated legumes in the field. ACIAR Monograph. 1989; 2(7):76.
- Reed MG, Scott AD. Chemical extraction of potassium from soils and micaceous minerals with solution containing sodium tetraphenylboron, II. Biotite. *Soil Science Society of American Proceedings*. 1962; 26:41-45.
- Ruiz-Lozano JM, Del Mar Alguacil M, Barzana G, Vernieri P, Aroca R. Exogenous ABA accentuates the differences in root hydraulic properties between mycorrhizal and non mycorrhizal maize plants through regulation of PIP aquaporins. *Plant Molecular Biology*. 2009; 70:565-579.
- Subbiah K. Effect of N, P and biofertilizers on yield and nutrient uptake in chilli and bellary onion. *Madras Agric. J.* 1994; 81:277-279.
- Tinker PB. The role of microorganisms in mediating and facilitating the uptake of plant nutrients from soil. *Plant and Soil*. 1984; 76:77-91.
- Turk MA, Assaf TA, Hameed KM, Altawaha AM. Significance of mycorrhiza. *World Journal of Agricultural Sciences*. 2006; 2(1):16-20
- Witham FH, Blaydes DF, Devlin RM. Experiments in Plant Physiology. 1st Edn., Van Nostrand, New York, 1971, 44-46.
- Yadav BD, Khandelwal RB, Sharma YK. Use of bio-fertilizer (Azospirillum) in onion. *Indian Journal Horticulture*. 2005; 62(2):168-70.
- Yousefi AA, Khavazi K, Moezi AA, Rejali F, Nadian NH. Phosphate solubilizing bacteria and arbuscular mycorrhizal fungi impacts on inorganic phosphorus fractions and wheat growth. *World Applied Sciences Journal*. 2011; 15(9):1310-1318.
- Zehra Ekin 2010. Performance of phosphate solubilizing bacteria for improving growth and yield of sunflower (*Helianthus annuus* L.) in the presence of phosphorus fertilizer. *Af. J. Biotech.* 9: 3794-00.