



STANDARDIZATION OF SEED TESTING PROTOCOL IN GRAIN AMARANTH (*Amaranthus hypochondriacus* L.)

Vijay, G.¹, Shashibhaskar M.S.^{2*}, Parashivamurthy¹, Madhusudan, K.³ and Anand S.R.⁴

¹ Department of Seed Science and Technology, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, India

² University Examination Centre, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, India

³ Registrar, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, India

⁴ AICRN on Weed Management, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, India

*Corresponding author E-mail: shashibhaskarms@gmail.com

(Date of Receiving-05-07-2026; Date of Revision-25-08-2026; Date of Acceptance-03-09-2026)

ABSTRACT

A laboratory experiment was conducted at Department of Seed Science and Technology, College of Agriculture, UAS, GKVK, Bengaluru. Grain amaranth seeds were subjected to germination at different substrates (sand, between paper and top of paper) and temperatures (20, 25, 30, 20/25 and 25/30 °C). Among the substrates, higher seed germination (80.06 %), mean seedling length (8.90 cm), mean seedling dry weight (4.25 mg), speed of germination (31.40) and seed vigour index-I and II (720 and 344) were recorded in between paper method. Among the temperatures, higher germination (86.55 %), mean seedling length (10.20 cm), mean seedling dry weight (4.96 mg) and seed vigour index-I and II (884 and 430) were recorded at alternative temperature of 25/30 °C.

Key words: Seed quality, substrate, temperature, seed testing.

Introduction

Grain amaranths belong to Amaranthaceae family and are herbaceous annual crop plants. They grow rapidly in a wide range of agroecological environments ranging from tropics to semi-arid and arid areas (Alemayehu *et al.*, 2015). Grain amaranth is a promising pseudo cereal crop to feed the human population (Mekonnen *et al.*, 2018) and a good forage for livestock (Mustafa *et al.*, 2011; Ahrar *et al.*, 2020). It has pale seeds and is known for its high nutritional value, being particularly rich in lysine (5%) and sulphur amino acids (4.4%). These amino acids are essential for a well-balanced human diet and are typically limited in the plant kingdom. Grain amaranth is often referred to as the “Poor man’s spinach” due to its affordability and is recommended as a means to combat malnutrition in various parts of the world. Grain amaranth surpasses cereal crops in terms of protein quality, primarily due to its high content of essential amino acids, including lysine, methionine and cysteine. Its significance lies in the excellent quality of its leaves and tender stems

as a food source, along with its high biocompatibility, nutritional value, antioxidants, mineral content, and cost-effectiveness compared to cereal crops (FAO, 2020). The nutritional attributes of grain amaranth, such as fiber and iron content, as well as its low levels of saturated fats and gluten-free nature, became widely recognized. Consequently, the crop has gained importance and experienced a resurgence as a health-conscious crop in numerous countries, including India.

One notable advantage of grain amaranth is its possession of the C4 pathway, which allows for efficient photosynthesis rates. This crop exhibits remarkable adaptability, capable of thriving in harsh environments. The genus amaranth is characterized by dibasic chromosomes, with a distribution nearly equal across the amaranth section and a chromosomal makeup of ($x = 16, 17$). Grant (1959) has documented the chromosome numbers of 30 *Amaranthus* species. Among these, four species have $2n = 32$ chromosomes, while the remaining species have $2n = 34$. Within the 30 species, *Amaranthus*

Table 1: Effect of substrate and temperature on seed germination and mean seedling length of grain amaranth.

Treatments	Seed germination (%)						Mean seedling length (cm)					
	T ₁	T ₂	T ₃	T ₄	T ₅	Mean	T ₁	T ₂	T ₃	T ₄	T ₅	Mean
S ₁	63.33	81.33	72.33	77.33	84.67	75.80	7.00	8.93	7.80	8.40	9.77	8.38
S ₂	71.67	83.33	75.33	80.33	89.67	80.06	7.50	9.53	8.13	8.73	10.60	8.90
S ₃	69.33	82.33	73.33	79.33	85.33	77.93	7.33	9.30	8.00	8.60	10.23	8.69
Mean	68.11	82.33	73.66	79.00	86.55		7.27	9.25	7.97	8.57	10.20	
	S		T		S×T		S		T		S×T	
SEm±	0.61		0.78		1.36		0.05		0.06		0.11	
CD (P=0.05)	1.76		2.27		NS		0.14		0.18		NS	
CV (%)	3.03						2.23					
*NS: Non-significant, S ₁ : Top of paper (TP), S ₂ : Between paper (BP), S ₃ : Sand, T ₁ : 20°C, T ₂ : 25°C, T ₃ : 30°C, T ₄ : 20/25°C and T ₅ : 25/30°C												

caudatus L. (2n = 32), *Amaranthus cruentus* L. (2n = 34), and *Amaranthus hypochondriacus* L. (2n = 32) are domesticated species. Among these three, *Amaranthus caudatus* L. exhibits a drooping growth habit, while the other two species are erect.

The use of quality seed is important for successful crop production. Therefore, it is crucial to increase the seed production of grain amaranth to meet growing demand. The quality of the seed lot is primarily evaluated through a seed testing procedure. Germination test has been accepted as an index of seed quality, which depends on the intrinsic characteristics of seeds and environmental factors such as temperature and substrate. Temperature affects the rate of imbibition and chemical reactions during germination process which influences uniformity and overall germination rate. Some species respond better at constant temperature while others best germinate at alternate temperature. The germination medium used plays a crucial role in the germination process of seeds, as it must meet specific requirements such as substrate to be non-toxic, free from moulds, and provide sufficient aeration and moisture for the seeds to germinate successfully. Before recommending a particular temperature and medium for adoption, it is necessary to standardize and determine the optimal temperature and

medium for testing the germination potential of grain amaranth. Currently no standardized seed testing protocol have been developed for grain amaranth (*Amaranthus hypochondriacus* L.). Therefore, the current study aims at developing methodology to evaluate germination and to establish a standard for routine in grain amaranth seed quality assessment.

Material and Methods

The present investigation was carried out in the Department of Seed Science and Technology, University of Agricultural Sciences, GKV, Bengaluru. The seeds of grain amaranth variety KBGA-15 were collected from AICRP on Potential Crop, UAS, GKV, Bengaluru. For seed germination testing protocol, standard growing media viz., top of paper, between paper and sand were used and temperature viz., constant temperature of 20°C, 25°C and 30°C and alternative temperature of 20/25°C and 25/30°C were employed in the experiment for standardization of substrate and temperature regime. The seeds were put to germination in top of paper by using blotters, between paper and trays containing sand. The germination test was carried out in walk in seed germinator. For alternate temperature, the lower temperature level was maintained for 16 hours and the higher temperature for 8 hours in a cycle of 24 hours. Seed quality parameters viz., seed germination, mean seedling length, mean seedling dry weight, speed of germination, seed vigour index-I and II were evaluated.

Results and Discussion

Seed germination (%)

The data on seed germination (%) as influenced by different media with different temperatures is presented in Table 1 and Fig. 1. The seed germination percentage was significantly higher (80.06 %) in between paper method (S₂), followed by top of paper method (S₃) (77.93 %) and lower seed germination (75.80 %) was recorded in sand method (S₁). Higher seed germination obtained




Fig. 1: Effect of substrate and temperature on seed germination of grain amaranth. [S₁: Top of paper (TP), S₂: Between paper (BP), S₃: Sand, T₁: 20°C, T₂: 25°C, T₃: 30°C, T₄: 20/25°C and T₅: 25/30°C]

mg) and lower mean seedling dry weight (3.34 mg) was recorded in seeds tested under temperature of 20 °C (T_1). Lima *et al.*, (2011) concluded that alternate temperature 20/30 °C and 20/35 °C provided good dry mass of seedling for *Caesalpinia pyramidalis* tul seeds.

Speed of germination

The data on speed of germination as influenced by different media with different temperatures is presented in Table 2. Significantly maximum speed of germination (31.40) was recorded in between paper method (S_2) followed by top of paper method (S_3) (29.77) and lower speed of germination (27.73) was recorded in sand method (S_1). Seeds tested in temperature of 30 °C (T_3) recorded significantly higher speed of germination (38.94) followed by alternating temperature of 25/30 °C (T_5) (35.32) and lower speed of germination (19.35) was recorded in seeds tested under temperature of T_1 (20 °C). When seeds were exposed to low temperature (20 °C), in general, there was a delay in germination from reduced metabolic activities. On the contrary, when seeds were placed in between paper at high temperature (30 °C) seeds germinated more rapidly as a result of protein denaturation coming from increased metabolic activities. This result was similar to the findings of De Jesus *et al.*, (2017) in quinoa seeds.

Seed Vigour Index-I

The data on seed vigour index-I as influenced by different media with different temperatures is presented in Table 3. Significantly higher seed vigour index-I (720) was recorded in between paper method (S_2) followed by top of paper method (S_3) (684) and lower seed vigour index-I (542) was recorded in sand method (S_1). Seeds tested in alternating temperature of 25/30 °C (T_5) recorded significantly higher seed vigour index-I (884) followed by temperature of 25 °C (T_2) (763) and lower seed vigour index-I (497) was recorded in 20 °C (T_1). This result was similar to the findings Buriro *et al.*, (2011) in wheat seed.

Seed vigour index-II

The data on seed vigour index-II as influenced by different media with different temperatures is presented in Table 3. Significantly maximum seed vigour index-II (344) was recorded in between paper method (S_2) followed by top of paper method (S_3) (325) and minimum (303) was recorded in sand method (S_1). Seeds tested in alternating temperature of 25/30 °C (T_5) recorded significantly higher seed vigour index-II (430) followed by temperature of 25 °C (T_2) (364) and less (228) was recorded in T_1 (20 °C). Seed vigour indices are sign of fast germination and speed of growth. These results are

in lines with the results of Geetha (2011) for *Cenchrus glaucus*.

Conclusion

The experimental results revealed that both substrate and temperature significantly influenced the seed germination and seedling performance in grain amaranth. Amongst the substrate, the between-paper (BP) method recorded the superior performance in seed germination, mean seedling length, mean seedling dry weight, speed of germination, seed vigour index-I, and seed vigour index-II. Amongst the temperatures, germination and seedling performance were superior under the alternating temperature of 25/30°C, indicating that this temperature regime provides the most favourable conditions for seed germination and subsequent seedling growth. Therefore, it is concluded that the germination test of grain amaranth could be conducted at an alternative temperature of 25/30°C using between paper as a substratum.

References

- Ahrar, A., Paknejad F., Tabatabaei S.A., Aghayari F., Soltani E. (2020). Evaluation of forage Amaranth (*Amaranthus hypochondriacus* L.) yield via comparing drought tolerance and susceptibility indices. *Ital. J. Agrometeorol.*, **3**, 31–40.
- Alemayehu, F.R., Bendevis M.A. and Jacobsen S.E. (2015). The potential for utilizing the seed crop amaranth (*Amaranthus* spp.) in East Africa as an alternative crop to support food security and climate change mitigation. *J. Agro. Crop Sci.*, **201(5)**, 321–329.
- Alves, C.Z., Silva J.B. and Candido A.C.S. (2015). Methodology for conducting the germination test in guava seeds. *Rev. Cienc. Agron.*, **46(3)**, 615–621.
- Alves, E.U., Paula R.C., Oliveira A.P., Bruno R.L.A. and Diniz A.A. (2002). Germination of *Mimosa caesalpiniaefolia* Benth seeds on different substrates and temperatures, *Rev. Bras. Sementes*, **24(1)**, 169–178.
- Anonymous (2021). International Rules for Seed Testing (ISTA). *Seed Sci. Technol.*, **24**, 1–335.
- Buriro, M., Oad F.C., Keerio M.I., Tunio S., Gandahi A.W., Hassan S.W.U. and Oad S.M. (2011). Wheat seed germination under the influence of temperature regimes. *Sarhad J. Agric.*, **27(4)**, 539–543.
- De Jesus Souza, F.F., De Souza J.E.A., Souza N.O.S., Spehar C.R. and De Jesus T.F. (2017). Standardizing germination tests for quinoa seeds. *Afr. J. Agric. Res.*, **12(3)**, 155–160.
- FAO (2020). Fruits and vegetables-your dietary essentials, *Backgr. Pap.*, **2(5)**, 81.
- Geetha, R. (2011). Standardization of germination test requirements for *Cenchrus glaucus*. *Intl. J. Plant. Sci.*, **6(1)**, 161–163.
- Gowda, R., Sowmya K.J. and Sriram I. (2012). Protocol for germination testing in some important medicinal crops.

- Res. J. Agric. Sci.*, **3(6)**, 1251-54.
- Grant, W.F. (1959). Cytogenetic studies in *Amaranthus*. *Can. J. Genet. Cytol.*, **1(4)**, 313-328.
- Lima, C.R.D., Pacheco M.V., Bruno R.D.L.A., Ferrari C.D.S., Braga Junior J.M. and Bezerra A.K.D. (2011). Temperature and substrate effects on the germination of *Caesalpinia pyramidalis* tul. seeds. *Rev. Bras. Sementes*, **33(2)**, 216-222.
- Mekonnen, G., Woldesenbet M., Teshale T., Biru T. (2018). *Amaranthus caudatus* production and nutrition contents for food security and healthy living in Menit Shasha, Menit Goldya and Maji districts of bench Maji zone, southwestern Ethiopia. *Nutr. Food Sci. Int. Journal.*, **7(3)**, 001–007.
- Muhl, Q.E., Du Toit E.S. and Robbertse P.J. (2011). Temperature effect on seed germination and seedling growth of *Moringa oleifera* Lam. *Seed Sci. Technol.*, **39(1)**, 208-213.
- Mustafa, A.F., Seguin P. and Gelinas B. (2011). Chemical composition, dietary fibre, tannins and minerals of grain amaranth genotypes. *Int. J. Food Sci. Nutr.*, **62(7)**, 750–754.
- Oliveira, A.K.M., Souza S.A., Souza J.S. and Carvalho J.M.B. (2015). Temperature and substrate influences on seed germination and seedling formation in *callisthene fasciculata*. *Rev. Arvore*, **39(3)**, 487-495.